

Dover Books Explaining Science and Mathematics

- The Rise of the New Physics*, A. d'Abrio 2 vol. set \$1.00
- The Evolution of Scientific Thought from Newton to Einstein*, A. d'Abrio \$2.00
- Elementary Concepts of Topology*, Paul Alexandroff \$1.00
- Language, Truth and Logic*, A. J. Ayer \$1.25
- A Dover Science Sampler* (61 page anthology), ed. by G. Barkin FREE
- The Sky and Its Mysteries*, E. A. Beut Clothbound \$3.50
- The Story of X-rays*, A. Bleich \$1.35
- Space and Time*, Emile Borel \$1.45
- Experiment & Theory in Physics*, Max Born 75¢
- The Restless Universe*, Max Born \$2.00
- Arithmetical Excursions*, H. & J. Bowers \$1.65
- Soap Bubbles, Their Color and the Forces Which Mould Them*, G. F. Boys 95¢
- Concerning the Nature of Things*, Sir William Bragg \$1.35
- The Universe of Light*, Sir William Bragg \$1.85
- The Nature of Physical Theory*, P. W. Bridgman \$1.25
- Matter and Light: The New Physics*, Louis de Broglie \$1.60
- Foundations of Science: The Philosophy of Theory & Experiment*,
N. Campbell \$2.95
- What Is Science?*, N. R. Campbell \$1.25
- The Common Sense of the Exact Sciences*, W. K. Clifford \$1.60
- The Internal Constitution of the Stars*, Arthur S. Eddington \$2.25
- The Story of Atomic Theory & Atomic Energy*, I. G. Fernberg \$1.45
- Insect Life & Insect Natural History*, S. W. Frost \$2.25
- Fads & Fallacies in the Name of Science*, Martin Gardner \$1.50
- Freshwater Microscopy*, W. Garret Clothbound \$5.95
- The Psychology of Invention in the Mathematical Field*, J. Hadamard \$1.25
- Popular Scientific Essays*, H. Helmholtz \$1.75
- On the Sensations of Tone*, H. Helmholtz \$2.95
- The Strange Story of the Quantum*, Banesh Hoffmann \$1.45
- The Realm of the Nebulae*, E. Hubble \$1.50
- Satellites and Scientific Research*, D. King Hele Clothbound \$1.00
- Trigonometry Refresher for Technical Men*, A. Klaf \$2.00
- Calculus Refresher for Technical Men*, A. Klaf \$2.00
- An Introduction to Symbolic Logic*, Suzanne Langer \$1.75
- Foundations of Physics*, R. Lindley & H. Margenau \$2.45

Presented to

Ludion Institute of Astrophysics

by D. Wechter

April 1977

(RICHARD H. WECHTER

514.13

MAN.1

NON-EUCLIDEAN GEOMETRY

NON-EUCLIDEAN GEOMETRY

BY
HENRY PARKER MANNING

Late Professor of Mathematics

Brown University

DOVER PUBLICATIONS, INC.

NEW YORK

IIA Lib.,

Published in the United Kingdom by Constable
and Company Limited, 10 Orange Street, London
W. C. 2.

This new Dover edition, first published in 1963,
is an unabridged and unaltered republication of
the work first published by Ginn and Company
in 1901.

Manufactured in the United States of America

Dover Publications, Inc.
180 Varick Street
New York 14, N. Y.

PREFACE

NON-EUCLIDEAN Geometry is now recognized as an important branch of Mathematics. Those who teach Geometry should have some knowledge of this subject, and all who are interested in Mathematics will find much to stimulate them and much for them to enjoy in the novel results and views that it presents.

This book is an attempt to give a simple and direct account of the Non-Euclidean Geometry, and one which presupposes but little knowledge of Mathematics. The first three chapters assume a knowledge of only Plane and Solid Geometry and Trigonometry, and the entire book can be read by one who has taken the mathematical courses commonly given in our colleges.

No special claim to originality can be made for what is published here. The propositions have long been established, and in various ways. Some of the proofs may be new, but others, as already given by writers on this subject, could not be improved. These have come to me chiefly through the translations of Professor George Bruce Halsted of the University of Texas.

I am particularly indebted to my friend, Arnold B. Chace, Sc.D., of Valley Falls, R. I., with whom I have studied and discussed the subject.

HENRY P. MANNING.

PROVIDENCE, January, 1901.

CONTENTS

	PAGE
INTRODUCTION	1

CHAPTER I

PANGEOMETRY

I. PROPOSITIONS DEPENDING ONLY ON THE PRINCIPLE OF SUPERPOSITION	3
II. PROPOSITIONS WHICH ARE TRUE FOR RESTRICTED FIGURES .	6
III. THE THREE HYPOTHESES	11

CHAPTER II

THE HYPERBOLIC GEOMETRY

I. PARALLEL LINES	31
II. BOUNDARY-CURVES AND SURFACES, AND EQUIDISTANT-CURVES AND SURFACES	43
III. TRIGONOMETRICAL FORMULÆ	53

CHAPTER III

THE ELLIPTIC GEOMETRY	62
---------------------------------	----

CHAPTER IV

ANALYTIC NON-EUCLIDEAN GEOMETRY

I. HYPERBOLIC ANALYTIC GEOMETRY	69
II. ELLIPTIC ANALYTIC GEOMETRY	80
III. ELLIPTIC SOLID ANALYTIC GEOMETRY	86
HISTORICAL NOTE	91

NON-EUCLIDEAN GEOMETRY

NON-EUCLIDEAN GEOMETRY

INTRODUCTION

THE axioms of Geometry were formerly regarded as laws of thought which an intelligent mind could neither deny nor investigate. Not only were the axioms to which we have been accustomed found to agree with our experience, but it was believed that we could not reason on the supposition that any of them are not true. It has been shown, however, that it is possible to take a set of axioms, wholly or in part contradicting those of Euclid, and build up a Geometry as consistent as his.

We shall give the two most important Non-Euclidean Geometries.* In these the axioms and definitions are taken as in Euclid, with the exception of those relating to parallel lines. Omitting the axiom on parallels,† we are led to three hypotheses; one of these establishes the Geometry of Euclid, while each of the other two gives us a series of propositions both interesting and useful. Indeed, as long as we can examine but a limited portion of the universe, it is not possible to prove that the system of Euclid is true, rather than one of the two Non-Euclidean Geometries which we are about to describe.

We shall adopt an arrangement which enables us to prove first the propositions common to the three Geometries, then to produce a series of propositions and the trigonometrical formulæ for each of the two Geometries which differ from

* See Historical Note, p. 93.

† See p. 91.

that of Euclid, and by analytical methods to derive some of their most striking properties.

We do not propose to investigate directly the foundations of Geometry, nor even to point out all of the assumptions which have been made, consciously or unconsciously, in this study. Leaving undisturbed that which these Geometries have in common, we are free to fix our attention upon their differences. By a concrete exposition it may be possible to learn more of the nature of Geometry than from abstract theory alone.

Thus we shall employ most of the terms of Geometry without repeating the definitions given in our text-books, and assume that the figures defined by these terms exist. In particular we assume :

I. The existence of straight lines determined by any two points, and that the shortest path between two points is a straight line.

II. The existence of planes determined by any three points not in a straight line, and that a straight line joining any two points of a plane lies wholly in the plane.

III. That geometrical figures can be moved about without changing their shape or size.

IV. That a point moving along a line from one position to another passes through every point of the line between, and that a geometrical magnitude, for example, an angle, or the length of a portion of a line, varying from one value to another, passes through all intermediate values.

In some of the propositions the proof will be omitted or only the method of proof suggested, where the details can be supplied from our common text-books.

CHAPTER I

PANGEOMETRY

I. PROPOSITIONS DEPENDING ONLY ON THE PRINCIPLE OF SUPERPOSITION

1. Theorem. *If one straight line meets another, the sum of the adjacent angles formed is equal to two right angles.*


2. Theorem. *If two straight lines intersect, the vertical angles are equal.*

3. Theorem. *Two triangles are equal if they have a side and two adjacent angles, or two sides and the included angle, of one equal, respectively, to the corresponding parts of the other.*

4. Theorem. *In an isosceles triangle the angles opposite the equal sides are equal.*

Bisect the angle at the vertex and use (3).

5. Theorem. *The perpendiculars erected at the middle points of the sides of a triangle meet in a point if two of them meet, and this point is the centre of a circle that can be drawn through the three vertices of the triangle.*



Proof. Suppose EO and FO meet at O . The triangles AFO and BFO are equal by (3). Also, AEO and CEO are equal.


Hence, CO and BO are equal, being each equal to AO . The triangle BCO is, therefore, isosceles, and OD if drawn bisecting the angle BOC will be perpendicular to BC at its middle point.

6. Theorem. *In a circle the radius bisecting an angle at the centre is perpendicular to the chord which subtends the angle and bisects this chord.*

7. Theorem. *Angles at the centre of a circle are proportional to the intercepted arcs and may be measured by them.*


8. Theorem. *From any point without a line a perpendicular to the line can be drawn.*

Proof. Let P' be the position which P would take if the plane were revolved about AB into coincidence with itself. The straight line PP' is then perpendicular to AB .



9. Theorem. *If oblique lines drawn from a point in a perpendicular to a line cut off equal distances from the foot of the perpendicular, they are equal and make equal angles with the line and with the perpendicular.*


10. Theorem. *If two lines cut a third at the same angle,*



that is, so that corresponding angles are equal, a line can be drawn that is perpendicular to both.

Proof. Let the angles FMB and MND be equal, and through H , the middle point of MN , draw LK perpendicular to CD ; then LK will also be perpendicular to AB . For the two triangles LMH and KNH are equal by (3).

11. Theorem. *If two equal lines in a plane are erected perpendicular to a given line, the line joining their extremities makes equal angles with them and is bisected at right angles by a third perpendicular erected midway between them.*



Let AC and BD be perpendicular to AB , and suppose AC and BD equal. The angles at C and D made with a line joining these two points are equal, and the perpendicular HK erected at the middle point of AB is perpendicular to CD at its middle point.


Proved by superposition.

12. Theorem. *Given as in the last proposition two perpendiculars and a third perpendicular erected midway between them; any line cutting this third perpendicular at right angles, if it cuts the first two at all, will cut off equal lengths on them and make equal angles with them.*

Proved by superposition.

Corollary. *The last two propositions hold true if the angles at A and B are equal acute or equal obtuse angles, HK being*


perpendicular to AB at its middle point. If $AC = BD$, the angles at C and D are equal, and HK is perpendicular to CD at its middle point; or, if CD is perpendicular to HK



at any point, K , and intersects AC and BD , it will cut off equal distances on these two lines and make equal angles with them.

II. PROPOSITIONS WHICH ARE TRUE FOR RESTRICTED FIGURES

The following propositions are true at least for figures whose lines do not exceed a certain length. That is, if there is any exception, it is in a case where we cannot apply the theorem or some step of the proof on account of the length of some of the lines. For convenience we shall use the word *restricted* in this sense and say that a theorem is true for restricted figures or in any restricted portion of the plane.




1. **Theorem.** *The exterior angle of a triangle is greater than either opposite interior angle (Euclid, I, 16).*

Proof. Draw AD from A to the middle point of the opposite side and produce it to E , making $DE = AD$. The two triangles ADC and EBD are equal, and the angle FBD , being greater than the angle EBD , is greater than C .

Corollary. *At least two angles of a triangle are acute.*

2. Theorem. *If two angles of a triangle are equal, the opposite sides are equal and the triangle is isosceles.*



Proof. The perpendicular erected at the middle point of the base divides the triangle into two figures which may be made to coincide and are equal. This perpendicular, therefore, passes through the vertex, and the two sides opposite the equal angles of the triangle are equal.


3. Theorem. *In a triangle with unequal angles the side opposite the greater of two angles is greater than the side opposite the smaller; and conversely, if the sides of a triangle are unequal the opposite angles are unequal, and the greater angle lies opposite the greater side.*

4. Theorem. *If two triangles have two sides of one equal, respectively, to two sides of the other, but the included angle of the first greater than the included angle of the second, the third side of the first is greater than the third side of the second; and conversely, if two triangles have two sides of*

one equal, respectively, to two sides of the other, but the third side of the first greater than the third side of the second, the angle opposite the third side of the first is greater than the angle opposite the third side of the second.

5. Theorem. *The sum of two lines drawn from any point to the extremities of a straight line is greater than the sum of two lines similarly drawn but included by them.*

6. Theorem. *Through any point one perpendicular only can be drawn to a straight line.*



Proof. Let P' be the position which P would take if the plane were revolved about AB into coincidence with itself. If we could have two perpendiculars, PC and PD , from P to AB , then CP' and DP' would be continuations of these lines and we should have two different straight lines joining P and P' , which is impossible.

Corollary. *Two right triangles are equal when the hypotenuse and an acute angle of one are equal, respectively, to the hypotenuse and an acute angle of the other.*

7. Theorem. *The perpendicular is the shortest line that can be drawn from a point to a straight line.*

Corollary. *In a right triangle the hypotenuse is greater than either of the two sides about the right angle.*


8. Theorem. *If oblique lines drawn from a point in a perpendicular to a line cut off unequal distances from the foot of the perpendicular, they are unequal, and the more remote is the greater; and conversely, if two oblique lines drawn from a point in a perpendicular are unequal, the greater cuts off a greater distance from the foot of the perpendicular.*

9. Theorem. *If a perpendicular is erected at the middle point of a straight line, any point not in the perpendicular is nearer that extremity of the line which is on the same side of the perpendicular.*

Corollary. *Two points equidistant from the extremities of a straight line determine a perpendicular to the line at its middle point.*

10. Theorem. *Two triangles are equal when they have three sides of one equal, respectively, to three sides of the other.*


11. Theorem. *If two lines in a plane erected perpendicular to a third are unequal, the line joining their extremities makes unequal angles with them, the greater angle with the shorter perpendicular.*



Proof. Suppose $AC > BD$. Produce BD , making $BE = AC$. Then $BEC = ACE$. But $BDC > BEC$, by (1), and ACD is a part of ACE . Therefore, all the more $BDC > ACD$.

12. Theorem. *If the two angles at C and D are equal, the perpendiculars are equal, and if the angles are unequal, the perpendiculars are unequal, and the longer perpendicular makes the smaller angle.*

13. Theorem. *If two lines are perpendicular to a third, points on either equidistant from the third are equidistant from the other.*



Proof. Let AB and CD be perpendicular to HK , and on CD take any two points, C and D , equidistant from K ; then C and D will be equidistant from AB . For by superposition we can make D fall on C , and then DB will coincide with CA by (6).

The following propositions of Solid Geometry depend directly on the preceding and hold true at least for any restricted portion of space.

14. Theorem. *If a line is perpendicular to two intersecting lines at their intersection, it is perpendicular to all lines of their plane passing through this point.*

15. Theorem. *If two planes are perpendicular, a line drawn in one perpendicular to their intersection is perpendicular to the other, and a line drawn through any point of one perpendicular to the other lies entirely in the first.*


16. Theorem. *If a line is perpendicular to a plane, any plane through that line is perpendicular to the plane.*

17. Theorem. *If a plane is perpendicular to each of two intersecting planes, it is perpendicular to their intersection.*

III. THE THREE HYPOTHESES

The angles at the extremities of two equal perpendiculars are either right angles, acute angles, or obtuse angles, at least for restricted figures. We shall distinguish the three cases by speaking of them as the hypothesis of the right angle, the hypothesis of the acute angle, and the hypothesis of the obtuse angle, respectively.

1. Theorem. *The line joining the extremities of two equal perpendiculars is, at least for any restricted portion of the plane, equal to, greater than, or less than the line joining their feet in the three hypotheses, respectively.*




Proof. Let AC and BD be the two equal perpendiculars and HK a third perpendicular erected at the middle point of AB . Then HA and KC are perpendicular to HK , and KC is equal to, greater than, or less than HA , according as the angle at C is equal to, less than, or greater than the angle at A (II, 12). Hence, CD , the double of KC , is equal to, greater than, or less than AB in the three hypotheses, respectively.

Conversely, if CD is given equal to, greater than, or less than AB , there is established for this figure the first, second, or third hypothesis, respectively.

Corollary. *If a quadrilateral has three right angles, the sides adjacent to the fourth angle are equal to, greater than, or less than the sides opposite them, according as the fourth angle is right, acute, or obtuse.*

2. Theorem. *If the hypothesis of a right angle is true in a single case in any restricted portion of the plane, it holds true in every case and throughout the entire plane.*




Proof. We have now a rectangle; that is, a quadrilateral with four right angles. By the corollary to the last proposition, its opposite sides are equal. Equal rectangles can be placed together so as to form a rectangle whose sides shall be any given multiples of the corresponding sides of the given rectangle.

Now let $A'B'$ be any given line and $A'C'$ and $B'D'$ two equal lines perpendicular to $A'B'$ at its extremities. Divide $A'C'$, if necessary, into a number of equal parts so that one of these parts shall be less than AC , and on AC and BD lay off AM and BN equal to one of these parts, and draw MN . $ABNM$ is a rectangle; for otherwise MN would be greater than or

less than AB and CD , and the angles at M and N would all be acute angles or all obtuse angles, which is impossible, since their sum is exactly four right angles. Again, divide $A'B'$ into a sufficient number of equal parts, lay off one of these parts on AB and on MN , and form the rectangle $APQM$. Rectangles equal to this can be placed together so as exactly to cover the figure $A'B'D'C'$, which must therefore itself be a rectangle.

3. Theorem. *If the hypothesis of the acute angle or the hypothesis of the obtuse angle holds true in a single case within a restricted portion of the plane, the same hypothesis holds true for every case within any such portion of the plane.*



Proof. Let CD move along AC and BD , always cutting off equal distances on these two lines; or, again, let AC and BD move along on the line AB towards HK or away from HK , always remaining perpendicular to AB and their feet always at equal distances from H . The angles at C and D vary continuously and must therefore remain acute or obtuse, as the case may be, or at some point become right angles. There would then be established the hypothesis of the right angle, and the hypothesis of the acute angle or of the obtuse angle could not exist even in the single case supposed.

The angles at C and D could not become zero nor 180° in a restricted portion of the plane; for then the three lines AC , CD , and BD would be one and the same straight line.

4. Theorem. *The sum of the angles of a triangle, at least in any restricted portion of the plane, is equal to, less than, or greater than two right angles, in the three hypotheses, respectively.*




FIG. 1.




FIG. 2.


Proof. Given any right triangle, ABD (Fig. 1), with right angle at B , draw AC perpendicular to AB and equal to BD . In the triangles ADC and DAB , $AC = BD$ and AD is common, but DC is equal to, greater than, or less than AB in the three hypotheses, respectively. Therefore, $\angle DAC$ is equal to, greater than, or less than $\angle ADB$ in the three hypotheses, respectively (II, 4). Adding $\angle BAD$ to both of these angles, we have $\angle ADB + \angle BAD$ equal to, less than, or greater than the right angle $\angle BAC$.

Now at least two angles of any restricted triangle are acute. The perpendicular, therefore, from the vertex of the third angle upon its opposite side will meet this side within the triangle and divide the triangle into two right triangles. Therefore, in any restricted triangle the sum of the angles is equal to, less than, or greater than two right angles in the three hypotheses, respectively.

We will call the amount by which the angle-sum of a triangle exceeds two right angles its excess. The excess of a polygon of n sides is the amount by which the sum of its angles exceeds $n - 2$ times two right angles.

It will not change the excess if we count as additional vertices any number of points on the sides, adding to the sum of the angles two right angles for each of these points.

5. Theorem. *The excess of a polygon is equal to the sum of the excesses of any system of triangles into which it may be divided.*



Proof. If we divide a polygon into two polygons by a straight or broken line, we may assume that the two points where it meets the boundary are vertices. If the dividing line is a broken line, broken at p points, the total sum of the angles of the two polygons so formed will be equal to the sum of the angles of the original polygon plus four right angles for each of these p points, and the sides of the two polygons will be the sides of the original polygon, together with the $p + 1$ parts into which the dividing line is separated by the p points, each part counted twice.

Let S be the sum of the angles of the original polygon, and n the number of its sides. Let S' and n' , S'' and n'' have the same meanings for the two polygons into which it is divided. Then we have, writing R for right angle,

$$S' + S'' = S + 4pR,$$

and

$$n' + n'' = n + 2(p + 1).$$

Therefore, $S' - 2(n' - 2)R + S'' - 2(n'' - 2)R$


$$= S + 4pR - 2(n + 2p - 2)R$$

$$= S - 2(n - 2)R.$$

Any system of triangles into which a polygon may be divided is produced by a sufficient number of repetitions of the above process. Always the excess of the polygon is equal to the sum of the excesses of the parts into which it is divided.

We may extend the notion of excess and apply it to any combination of different portions of the plane bounded completely by straight lines.

Instead of considering the sum of the angles of a polygon, we may take the sum of the exterior angles. The amount by which this sum falls short of four right angles equals the excess of the polygon. We may speak of it as the deficiency of the exterior angles.



The sum of the exterior angles is the amount by which we turn in going completely around the figure, turning at each vertex from one side to the next. If we are considering a combination of two or more polygons, we must traverse the entire boundary and so as always to have the area considered on one side, say on the left.

6. Theorem. *The excess of polygons is always zero, always negative, or always positive.*


Proof. We know that this theorem is true of restricted triangles, but any finite polygon may be divided into a finite number of such triangles, and by the last theorem the excess of the polygon is equal to the sum of the excesses of the triangles.

When the excess is negative, we may call it deficiency, or speak of the excess of the exterior angles.

Corollary. *The excess of a polygon is numerically greater than the excess of any part which may be cut off from it by straight lines, except in the first hypothesis, when it is zero.*

The following theorems apply to the second and third hypotheses.

7. Theorem. *By diminishing the sides of a triangle, or even one side while the other two remain less than some fixed length, we can diminish its area indefinitely, and the sum of its angles will approach two right angles as limit.*




Proof. Let $ABDC$ be a quadrilateral with three right angles, A , B , and C . A perpendicular moving along AB will constantly increase or decrease; for if it could increase a part of the way and decrease a part of the way there would be different positions where the perpendiculars have the same length; a perpendicular midway between them would be perpendicular to CD also, and we should have a rectangle.

Divide AB into n equal parts, and draw perpendiculars through the points of division. The quadrilateral is divided into n smaller quadrilaterals, which can be applied one to another, having a side and two adjacent right angles the same in all. Beginning at the end where the perpendicular is the shortest, each quadrilateral can be placed entirely within the next. Therefore, the first has its area less than $\frac{1}{n}$ th of the area of the original quadrilateral, and its deficiency or excess less than $\frac{1}{n}$ th of the deficiency or excess of the whole. Now any triangle whose sides are all less than AC or BD , and one of whose sides is less than one of the subdivisions of AB ,

can be placed entirely within this smallest quadrilateral. Such a triangle has its area and its deficiency or excess less than $\frac{1}{n}$ th of the area and of the deficiency or excess of the original quadrilateral.

Thus, a triangle has its area and deficiency or excess less than any assigned area and deficiency or excess, however small, if at least one side is taken sufficiently small, the other two sides not being indefinitely large.

8. Theorem. *Two triangles having the same deficiency or excess have the same area.*



Proof. Let AOB and $A'OB'$ have the same deficiency or excess and an angle of one equal to an angle of the other. If we place them together so that the equal angles coincide, the triangles will coincide and be entirely equal, or there will be a quadrilateral common to the two, and, besides this, two smaller triangles having an angle the same in both and the same deficiency or excess. Putting these together, we find again a quadrilateral common to both and a third pair of triangles having an angle the same in both and the same deficiency or excess. We may continue this process indefinitely; unless we come to a pair of triangles which coincide; for at no time can one triangle of a pair be contained entirely within the other, since they have the same deficiency or excess.


Let so denote the sum of the sides opposite the equal angles of the first two triangles, sa the sum of the adjacent sides, and $s'a$ that portion of the adjacent sides counted twice, which is common to the two triangles when they are placed together. Writing o' and a' for the second pair of triangles, o'' and a'' for the third pair, etc., we have

$$\begin{aligned} sa &= s'a + so', & so &= sa', \\ sa' &= s'a' + so'', & so' &= sa'', \\ sa'' &= s'a'' + so''', \text{ etc.} & so'' &= sa''', \text{ etc.} \end{aligned}$$

$$\therefore sa = s'a + s'a'' + s'a'' + \dots,$$

$$sa' = s'a' + s'a''' + s'a'' + \dots$$


Therefore, the expressions $s'a$, $s'a'$, $s'a''$, \dots diminish indefinitely. Each of these is made up of a side counted twice from one and a side counted twice from the other of a pair of triangles. Thus, if we carry the process sufficiently far, the remaining triangles can be made to have at least one side as small as we please, while all the sides diminish and are less, for example, than the longest of the sides of the original triangles. Therefore, the areas of the remaining triangles diminish indefinitely, and as the difference of the areas remains the same for each pair of triangles, this difference must be zero. The triangles of each pair and, in particular, the first two triangles have the same area.



Let ABC and DEF have the same deficiency or excess, and suppose $AC < DF$. Produce AC to C' , making $AC' = DF$.

Then there is some point, B' , on AB between A and B such that $AB'C'$ has the same deficiency or excess and the same area as ABC . Place $AB'C'$ upon DEF so that AC' will coincide with DF , and let $DE'F$ be the position which it takes. If the triangles do not coincide, the vertex of each opposite the common side DF lies outside of the other. The two triangles have in common a triangle, say DOF , and besides this there remain of the two triangles two smaller triangles which have one angle the same in both and the same deficiency or excess. These two triangles, and therefore the original triangles, have the same area.


9. Theorem. *The areas of any two triangles are proportional to their deficiencies or excesses.*



Proof. A triangle may be divided into n smaller triangles having equal deficiencies or excesses and equal areas by lines drawn from one vertex to points of the opposite side. Each of these triangles has for its deficiency or excess $\frac{1}{n}$ th of the deficiency or excess of the original triangle, and for its area $\frac{1}{n}$ th of the area of the original triangle.

When the deficiencies or excesses of two triangles are commensurable, say in the ratio $m:n$, we can divide them into m and n smaller triangles, respectively, all having the same deficiency or excess and the same area. The areas of the given triangles will therefore be in the same ratio, $m:n$.

When the deficiencies or excesses of two triangles, A and B , are not commensurable, we may divide one triangle, A , as above, into any number of equivalent parts, and take parts equivalent to one of these as many times as possible from the



other, leaving a remainder which has a deficiency or excess less than the deficiency or excess of one of these parts. The portion taken from the second triangle forms a triangle, B' . A and B' have their areas proportional to their deficiencies or excesses, these being commensurable. Now increase indefinitely the number of parts into which A is divided. These parts will diminish indefinitely, and the remainder when we take B' from B will diminish indefinitely. The deficiency or excess and the area of B' will approach those of B , and the triangles A and B have their areas and their deficiencies or excesses proportional.

Corollary. *The areas of two polygons are to each other as their deficiencies or excesses.*

10. Theorem. *Given a right triangle with a fixed angle; if the sides of the triangle diminish indefinitely, the ratio of the opposite side to the hypotenuse and the ratio of the adjacent side to the hypotenuse approach as limits the sine and cosine of this angle.*

Proof. Lay off on the hypotenuse any number of equal lengths. Through the points of division A_1, A_2, \dots draw perpendiculars A_1C_1, A_2C_2, \dots to the base, and to these lines


produced draw perpendiculars A_2D_1 , A_3D_2 , \dots each from the next point of division of the hypotenuse.

The triangles OA_1C_1 and $A_2A_1D_1$ are equal (II, 6, Cor.).

$$C_2A_2 \geq C_1D_1 \quad \text{and} \quad C_1C_2 \leq D_1A_2;$$

therefore,
$$\frac{C_2A_2}{OA_2} \geq \frac{C_1A_1}{OA_1} \quad \text{and} \quad \frac{OC_2}{OA_2} \leq \frac{OC_1}{OA_1},$$

the upper sign being for the second hypothesis and the lower sign for the third hypothesis.



Assume
$$\frac{C_{r-1}A_{r-1}}{OA_{r-1}} \geq \frac{C_{r-2}A_{r-2}}{OA_{r-2}} \geq \dots \geq \frac{C_1A_1}{OA_1},$$

and
$$\frac{OC_{r-1}}{OA_{r-1}} \leq \frac{OC_{r-2}}{OA_{r-2}} \leq \dots \leq \frac{OC_1}{OA_1}.$$

Since
$$OA_{r-1} = (r-1)OA_1,$$

and also
$$= (r-1)A_{r-1}A_r,$$

the inequalities
$$\frac{OC_1}{OA_1} \geq \frac{OC_{r-1}}{OA_{r-1}} \quad \text{and} \quad \frac{C_1A_1}{OA_1} \geq \frac{C_{r-1}A_{r-1}}{OA_{r-1}}$$

applied to the angle at A_{r-1} become

$$\frac{A_{r-1}D_{r-1}}{A_{r-1}A_r} \geq \frac{C_{r-1}A_{r-1}}{OA_{r-1}} \quad \text{and} \quad \frac{D_{r-1}A_r}{A_{r-1}A_r} \leq \frac{OC_{r-1}}{OA_{r-1}}.$$

The first of these two inequalities may be written

$$\frac{A_{r-1}D_{r-1}}{C_{r-1}A_{r-1}} \geq \frac{A_{r-1}A_r}{OA_{r-1}}.$$

Add 1 to both members,

$$\frac{C_{r-1}D_{r-1}}{C_{r-1}A_{r-1}} \geq \frac{OA_r}{OA_{r-1}},$$

or

$$\frac{C_{r-1}D_{r-1}}{OA_r} \geq \frac{C_{r-1}A_{r-1}}{OA_{r-1}}.$$

But

$$C_r A_r \geq C_{r-1} D_{r-1}.$$

\therefore

$$\frac{C_r A_r}{OA_r} \geq \frac{C_{r-1} A_{r-1}}{OA_{r-1}}.$$

Again,

$$C_{r-1} C_r \leq D_{r-1} A_r.$$

Hence, from the second inequality above, we have

$$\frac{C_{r-1} C_r}{A_{r-1} A_r} \leq \frac{OC_{r-1}}{OA_{r-1}},$$

or

$$\frac{C_{r-1} C_r}{OC_{r-1}} \leq \frac{A_{r-1} A_r}{OA_{r-1}}.$$

Add 1 to both members,

$$\frac{OC_r}{OC_{r-1}} \leq \frac{OA_r}{OA_{r-1}},$$


or

$$\frac{OC_r}{OA_r} \leq \frac{OC_{r-1}}{OA_{r-1}}.$$

The ratios $\frac{CA}{OA}$ and $\frac{OC}{OA}$ being less than 1, and always increasing or always decreasing when the hypotenuse decreases, approach definite limits. These limits are continuous functions of A ; if we vary the angle of any right triangle continuously, keeping the hypotenuse some fixed length, the other two sides will vary continuously, and the limits of their ratios to the hypotenuse must, therefore, vary continuously.

Calling the limits for the moment sA and cA , we may extend their definition, as in Trigonometry, to any angles, and prove that all the formulæ of the sine and cosine hold for these functions. Then for certain angles, 30° , 45° , 60° , we can prove

that they have the same values as the sine and cosine, and their values for all other angles as determined from their values for these angles will be the same as the corresponding values of the sine and cosine.



Draw a perpendicular, CF , from the right angle C to the hypotenuse AB . The angle FCB is not equal to A , but the difference, being proportional to the difference of areas of the two triangles ABC and FBC , diminishes indefinitely when the sides of the triangles diminish. From the relation

$$\frac{AF}{AC} \frac{AC}{AB} + \frac{FB}{BC} \frac{BC}{AB} = 1,$$

we have, by passing to the limit,

$$(cA)^2 + (sA)^2 = 1.$$

Let x and y be any two acute angles, and draw the figures used to prove the formulæ for the sine and cosine of the sum of two angles.

The angles x and y remaining fixed, we can imagine all of the lines to decrease indefinitely, and the functions sx , cx , sy , etc., are the limits of certain ratios of these lines.

$$\begin{aligned} \frac{CA}{OA} &= \frac{CE}{OB} \frac{OB}{OA} + \frac{EA}{BA} \frac{BA}{OA}, \\ \pm \frac{OC}{OA} &= \frac{OD}{OB} \frac{OB}{OA} - \frac{CD}{BA} \frac{BA}{OA} \end{aligned}$$



$$\left(-\frac{OC}{OA} \text{ in the second figure} \right).$$

The angles at M are equal in the two triangles EMB and CMO , and we may write

$$\frac{CM}{OM} = \frac{ME + \delta}{MB} = \frac{ME \pm CM + \delta}{MB \pm OM},$$

where δ has the limit zero.

$$\therefore \lim \frac{CE}{OB} = \lim \frac{CM}{OM} = sx.$$



The angle EAB , or x' , is not the same as x , but differs from x only by an amount which is proportional to the difference of the areas of the triangles OMC and MAB , and which, therefore, diminishes indefinitely. Thus, the limits of sx' and cx' are sx and cx .

Finally, as the two triangles ACN and BDN have the angle N in common, we may write

$$\frac{DN}{BN} = \frac{CN + \delta'}{AN} = \frac{CN - DN + \delta'}{AN - BN},$$

where the limit of δ' is zero.

$$\therefore \lim \frac{CD}{AB} = \lim \frac{CN}{AN} = sx.$$

Now at the limits our identities become

$$s(x + y) = sx \cdot cy + cx \cdot sy,$$

$$c(x + y) = cx \cdot cy - sx \cdot sy.$$

By induction, these formulæ are proved true for any angles. Other formulæ sufficient for calculating the values of these functions from their values for 30° , 45° , and 60° are obtained from these two by algebraic processes.

If the sides of an isosceles right triangle diminish indefinitely, the angle does not remain fixed but approaches 45° , and the ratios of the two sides to the hypotenuse approach as limits $s\,45^\circ$ and $c\,45^\circ$. Therefore, these latter are equal, and since the sum of their squares is 1, the value of each is $\frac{1}{\sqrt{2}}$, the same as the value of the sine and cosine of 45° .


Again, bisect an equilateral triangle and form a triangle in which the hypotenuse is twice one of the sides. When the sides diminish, preserving this relation, the angles approach 30° and 60° . Therefore, the functions, s and c , of these angles have values which are the same as the corresponding values of the sine and cosine of the same angles.

Corollary. *When any plane triangle diminishes indefinitely, the relations of the sides and angles approach those of the sides and angles of plane triangles in the ordinary geometry and trigonometry with which we are familiar.*

11. Theorem. *Spherical geometry is the same in the three hypotheses, and the formulæ of spherical trigonometry are exactly those of the ordinary spherical trigonometry.*

Proof. On a sphere, arcs of great circles are proportional to the angles which they subtend at the centre, and angles on a sphere are the same as the diedral angles formed by the planes of the great circles which are the sides of the angles. Their relations are established by drawing certain plane triangles which may be made as small as we please, and therefore may be assumed to be like the plane triangles in the hypothesis of a right angle. These relations are, therefore, those of the ordinary Spherical Trigonometry.

The three hypotheses give rise to three systems of Geometry, which are called the Parabolic, the Hyperbolic, and the Elliptic Geometries. They are also called the Geometries of Euclid, of Lobachevsky, and of Riemann. The following considerations exhibit some of their chief characteristics.



Given PC perpendicular to a line, CF ; on the latter we take

$$CD = PC,$$

$$DD' = PD,$$

$$D'D'' = PD', \text{ etc.}$$

Now if PC is sufficiently short (restricted), it is shorter than *any* other line from P to the line CF ; for any line as short as PC or shorter would be included in a restricted portion of the plane about the point P , for which the perpendicular is the shortest distance from the point to the line.

Therefore, $PD > PC$, $\therefore CD' > 2 CD$,
 $PD' > PC$, etc.; $CD'' > 3 CD$, etc.

Again, in the three hypotheses, respectively,

$$CPD \lesseqgtr \frac{\pi}{4}, \quad \text{and} \quad CDP \lesseqgtr \frac{\pi}{4},$$

$$DPD' \lesseqgtr \frac{1}{2} CPD, \quad CD'P \lesseqgtr \frac{1}{2} CDP,$$

$$D'PD'' \lesseqgtr \frac{1}{2} DPD', \text{ etc.}, \quad CD''P \lesseqgtr \frac{1}{2} CD'P, \text{ etc.}$$

At P we have a series of angles. In the first hypothesis there is an infinite number of these angles, and the series forms a geometrical progression of ratio $\frac{1}{2}$, whose value is

exactly $\frac{\pi}{2}$. In the second hypothesis there is also an infinite number of these angles, and the terms of the series are less than the terms of the geometrical progression. The value of the series is, therefore, less than $\frac{\pi}{2}$. In the third hypothesis we have a series whose terms are greater than those of the geometrical progression, and, therefore, whether the series is convergent or divergent, we can get more than $\frac{\pi}{2}$ by taking a sufficient number of terms. In other words, we can get a right angle or more than a right angle at P by repeating this process a certain finite number of times.

The angles at D, D', D'', \dots are exactly equal to the terms of the series of angles at P . In the first two hypotheses they approach zero as a limit.


The distances CD, CD', CD'', \dots increase each time by more than a definite quantity, CD ; therefore, if we repeat the process an unlimited number of times, these distances will increase beyond all limit. Thus, in the first and second hypotheses we prove that a straight line must be of infinite length.

In the hypothesis of the obtuse angle the line perpendicular to PC at the point P will intersect CF in a point at a certain finite distance from C , one of the D 's, or some point between. On the other side of PC this same perpendicular will intersect FC produced at the same distance. But we have assumed that two different straight lines cannot intersect in two points; therefore, for us the third hypothesis cannot be true unless the straight line is of finite length returning into itself, and these two points are one and the same point, its distance from C in either direction being one-half the entire length of the line. In this way, however, we can build up a consistent Geometry on the third hypothesis, and this Geometry it is which is called the Elliptic Geometry.

The constructions would have been the same, and very nearly all the statements would have been the same, if we had taken CD any arbitrary length on CF .

The restriction which we have placed upon some of the propositions of this chapter is necessary in the third hypothesis.


Thus, in the proof that the exterior angle of a triangle is greater than the opposite interior angle, the line AD drawn through the vertex A to the middle point D of the opposite



side was produced so as to make $AE = 2 AD$. If AD were greater than half the entire length of the straight line determined by A and D , this would bring the point E past the point A , and the angle CBE , which is equal to the angle C , instead of being a part of the exterior angle CBF , becomes greater than this exterior angle.

Again, if two angles of a triangle are equal and the side between them is just an entire straight line, it does not follow necessarily that the opposite sides are equal. It may be said,

however, that the opposite sides form one continuous line, and, therefore, this figure is not strictly a triangle, but a figure



somewhat like a lune. The points A and B are the same point, and the angles A and B are vertical angles.

Finally, though we assume that the shortest path between two points is a straight line, it is not always true that a straight line drawn between two points is the shortest path between them. We can pass from one point to another in two ways on a straight line; namely, over each of the two parts into which the two points divide the line determined by them. One of these parts will usually be shorter than the other, and the longer part will be longer than some paths along broken lines or curved lines.

When, however, the straight line is of infinite length, that is, in the hypothesis of the right angle and in the hypothesis of the acute angle, all the propositions of this chapter hold without restriction.

The Euclidean Geometry is familiar to all. We will now make a detailed study of the Geometry of Lobachevsky, and then take up in the same way the Elliptic Geometry.


CHAPTER II

THE HYPERBOLIC GEOMETRY

WE have now the hypothesis of the acute angle. Two lines in a plane perpendicular to a third diverge on either side of their common perpendicular. The sum of the angles of a triangle is less than two right angles, and the propositions of the last chapter hold without restriction.

I. PARALLEL LINES


From any point, P , draw a perpendicular, PC , to a given line, AB , and let PD be any other line from P meeting CB in D . If D move off indefinitely on CB , the line PD will approach a limiting position PE .



PE is said to be parallel to CB at P . PE makes with PC an angle, CPE , which is called the angle of parallelism for the perpendicular distance PC . It is less than a right angle by an amount which is the limit of the deficiency of the triangle PCD . On the other side of PC we can find another line parallel to CA and making with PC the same angle of parallelism. We say that PE is parallel to AB towards that part which is on the same side of PC with PE . Thus, at any

point there are two parallels to a line, but only one towards one part of the line. Lines through P which make with PC an angle greater than the angle of parallelism and less than its supplement do not meet AB at all. We write $\Pi(p)$ to denote the angle of parallelism for a perpendicular distance, p .

1. Theorem. *A straight line maintains its parallelism at all points.*



Let AB be parallel to CD at E and let F be any other point of AB on either side of E , to prove that AB is parallel to CD at F .

Proof. To H , on CD , draw EH and FH . If H move off indefinitely on CD , these two lines will approach positions of parallelism with CD . But the limiting position of EH is the line AB passing through F , and if the limiting position of FH were some other line, FK , F would be the limiting position of H , the intersection of EH and FH .

2. Theorem. *If one line is parallel to another, the second is parallel to the first.*


Given AB parallel to CD , to prove that CD is parallel to AB .

Proof. Draw AC perpendicular to CD . The angle CAB will be acute; therefore, the perpendicular CE from C to AB must fall on that side of A towards which the line AB is parallel to CD (Chap. I, II, 1). The angle ECD is then acute and less than CEB , which is a right angle. That is, we have

$$CAB < ACD, \text{ and } CEB > ECD.$$


If the line CE revolve about the point C to the position of CA , the angle at E will decrease to the angle A , and the angle at C will increase to a right angle. There will be some position, say CF , where these two angles become equal; that is,

$$CFB = FCD.$$



Draw MN perpendicular to CF at its middle point and revolve the figure about MN as an axis. CD will fall upon the original position of AB , and AB will fall upon the original position of CD . Therefore, CD is parallel to AB .

Corollary. FB and CD are both parallel to MN .




Proof. FB and CD are symmetrically situated with respect to MN , and cannot intersect MN since they do not intersect each other. Draw FH to H , on CD , intersecting MN in K . If H move off indefinitely on CD , FH will approach the position of FB as a limit. Now K cannot move off indefinitely before H does, for $FK < FH$. But again, when H moves off indefinitely, K cannot approach some limiting position at a

finite distance on MN ; for FB , and therefore CD , would then intersect MN and each other at this point. Therefore, H and K must move off together, and the limiting position of FH must be at the same time parallel to CD and MN .

In the same way we can prove that any line lying in a plane between two parallels must intersect one of them or be parallel to both.

3. Theorem. *Two lines parallel to a third towards the same part of the third are parallel to each other.*




First, when they are all in the same plane.

Let AB and EF be parallel to CD , to prove that they are parallel to each other.

Proof. Suppose AB lies between the other two. To H , any point on CD , draw AH and EH , and let K be the point where EH intersects AB . As H moves off indefinitely on CD , AH and EH approach as limiting positions AB and EF . Now K cannot move off indefinitely before H does, for $EK < EH$. But again, when H moves off indefinitely, K cannot approach some limiting position at a finite distance on AB ; for this point would be the intersection of AB and EF , and the limiting position of H , whereas H moves off indefinitely on CD . Therefore, H and K must move off together, and the limiting position of EH must be at the same time parallel to CD and AB .

If AB , lying between the other two, is given parallel to CD and EF , EF must be parallel to CD ; for a line through E parallel to CD would be parallel to AB , and only one line can be drawn through E parallel to AB towards the same part.



Second, when the lines are not all in the same plane.

Let AB and CD be two parallel lines and let E be any point not in their plane.

Proof. To H on CD draw AH and EH . As H moves off indefinitely, AH approaches the position of AB , and the plane EAH the position of the plane EAB . Therefore, the limiting position of EH is the intersection of the planes ECD and EAB . The intersection of these planes is, then, parallel to CD , and in the same way we prove that it is parallel to AB .


Now, if EF is given as parallel to one of these two lines towards the part towards which they are parallel, it must be the intersection of the two planes determined by them and the point E , and therefore parallel to the other line also.

4. Theorem. *Parallel lines continually approach each other.*

Let AB and CD be parallel, and from A and B , any points on AB , drop perpendiculars AC and BD to CD . Supposing that B lies beyond A in the direction of parallelism, we are to prove that $BD < AC$.


Proof. At H , the middle point of CD , erect a perpendicular meeting AB in K . The angle BKH is an acute angle, and the

angle AKH is an obtuse angle. Therefore, a perpendicular to HK at K must meet CA in some point, E , between C and A



and DB produced in some point, F , beyond B . But $DF = CE$ (Chap. I, I, 12); therefore, $DB < CA$.

Corollary. *If AB and CD are parallel and AC makes equal angles with them (like FC in 2 above), then EF , cutting off equal distances on these two lines, $AE = CF$, on the side towards which they are parallel, will be shorter than AC ,*



Proof. MN , perpendicular to AC at its middle point, is parallel to AB and bisects EF , the figure being symmetrical with respect to MN . EH , the half of EF , is less than AM , and therefore EF is less than AC .


5. Theorem. *As the perpendicular distance varies, starting from zero and increasing indefinitely, the angle of parallelism decreases from a right angle to zero.*

Proof. In the first place the angle of parallelism, which is acute as long as the perpendicular distance is positive, will be

made to differ from a right angle by less than any assigned value if we take a perpendicular distance sufficiently small.



For, ADE being any given angle as near a right angle as we please, we can take a point, L , on DE and draw LR perpendicular to DA at R . The angle RDL must increase to become the angle of parallelism for the perpendicular distance RD .



Now let p be the length of a given perpendicular PM , and let α be the amount by which its angle of parallelism differs from $\frac{\pi}{2}$; that is, say

$$\Pi(p) = \frac{\pi}{2} - \alpha.$$

PM , being perpendicular to MN , and H any point on MN , the angle MPH approaches as a limit the angle of parallelism, $\Pi(p)$, when H moves off indefinitely on MN . The line PH meets the line MN as long as $MPH < \Pi(p)$, and by taking MPH sufficiently near $\Pi(p)$, but less, we can make the angle MHP as small as we please (see p. 27).

In figure on page 38, let AC be perpendicular to AB , D being any point on AC and DE parallel to AB . Draw DK beyond DE , making with DE an angle, $EDK = \Pi(p)$, and make $DK = p$. TF , perpendicular to DK at K , will be parallel to DE and AB .

By placing PMN of the last figure upon DKT , we see that DC will meet KT in a point, G if

$$KDC < \Pi(p),$$

that is, if

$$ADE > 2\alpha.$$

Then in the right triangle DKG ,


$$DGK + KDG < \frac{\pi}{2}.$$

But

$$ADE + KDG = \frac{\pi}{2} + \alpha;$$

therefore,

$$DGK < ADE - \alpha.$$




Starting from the point G , we can repeat this construction, and each time we subtract from the angle of parallelism an amount greater than α . We can continue this process until the angle of parallelism becomes equal to or less than 2α .

If the point D move along AC , DE remaining constantly parallel to AB , the angle at D will constantly diminish, and by letting D move sufficiently far on AC we can reach a point where this angle becomes equal to or less than 2α .

Suppose D is at the point where the angle of parallelism is just 2α . Then, if we draw DK and TF as before, KT will be


parallel to DC . All the parallels to AB lying between AB and this position of TF meet AC , and as the parallel moves towards this position of TF , the angle of parallelism at D approaches zero, and the point D moves off indefinitely.



For an obtuse angle we may take p negative, and we have

$$\Pi(-p) = \pi - \Pi(p).$$

6. Theorem. *The perpendiculars erected at the middle points of the sides of a triangle are all parallel if two of them are parallel.*



Let A , B , and C be the vertices of the triangle, and D , E , and F , respectively, the middle points of the opposite sides. Suppose the perpendiculars at D and E are given parallel, to prove that the perpendicular at F is parallel to them.


Proof. Draw CM through C parallel to the two given parallel perpendiculars. CM forms with the two sides at C angles of parallelism $\Pi\left(\frac{a}{2}\right)$ and $\Pi\left(\frac{b}{2}\right)$, of which the angle at C is the sum or difference according as C lies between the given perpendiculars or on the same side of both. By properly diminishing these angles at C , keeping the lengths of CA and CB unchanged, we can make the perpendiculars at their middle points D and E intersect CM , and therefore each other, at any distance from C greater than $\frac{a}{2}$ and greater than $\frac{b}{2}$.

Let $A'B'C'$ be the triangle so formed, O the point where the two given perpendiculars meet, and $C'M'$ the line through O . In the triangle $A'B'C'$, the three perpendiculars meet at the point O (Chap. I, I, 5). Now we can let O move off on $C'M'$, the construction remaining the same. That is, we let the lines $C'A'$ and $C'B'$ rotate about C' without changing their lengths, in such a manner that the three perpendiculars $D'O$, $E'O$, and $F'O$ shall always pass through O . As O moves off indefinitely, the angles at C' approach $\Pi\left(\frac{a}{2}\right)$ and $\Pi\left(\frac{b}{2}\right)$ as limits, and the three perpendiculars approach positions of parallelism with $C'M'$ and with each other. But the triangle $A'B'C'$ approaches as a limit a triangle which is equal to ABC , having two sides and the included angle equal, respectively, to the corresponding parts of the latter. Therefore, in ABC the three perpendiculars are all parallel.

7. Theorem. *Lines which do not intersect and are not parallel have one and only one common perpendicular.*

Proof. Let AB and CD be the two lines, and from A , any point of AB , drop AC perpendicular to CD . If AC is not itself the common perpendicular, one of the angles which it makes with AB will be acute. Let this angle be on the side


towards AB , so that $BAC < \frac{\pi}{2}$. Draw AE parallel to CD on this same side of AC . The angle EAC is less than BAC , since AB is not parallel to CD and does not intersect it. Let AH be any line drawn in the angle EAC , intersecting CD at H . If H , starting from the position of C , move off indefinitely



on the line CD , the angle BAH will decrease from the magnitude of the angle BAC to the angle BAE . The angle AHC will decrease *indefinitely* from the magnitude of the angle at C , which is a right angle and greater than BAC . There will be some position for which $BAH = AHC$. In this position the line NM through the middle point of AH perpendicular to one of the two given lines will be perpendicular to the other, as proved in Chap. I, I, 10.

If there were two common perpendiculars we should have a rectangle, which is impossible in the Hyperbolic Geometry.

8. Theorem. *If the perpendiculars erected at the middle*





points of the sides of a triangle do not meet and are not parallel, they are all perpendicular to a certain line.

Proof. We can draw a line, AB , that will be perpendicular to two of these lines, and the perpendiculars from the three vertices of the triangle upon this line will be equal, by Chap. I, II, 13. A perpendicular to AB erected midway between any two of these three is perpendicular to the corresponding side of the triangle at its middle point (Chap. I, I, 11). Thus, all three of the perpendiculars erected at the middle points of the sides of the triangle are perpendicular to AB .

A line is parallel to a plane if it is parallel to its projection on the plane.

9. Theorem. *A line may be drawn perpendicular to a plane and parallel to any line not in the plane.*



Proof. Let AB be the given line and MN the plane. If AB meets the plane MN at a point, A , we take on its projection a length, AC , such that the angle at A equals $\Pi(AC)$. Then CD , perpendicular to the plane at C , will be parallel to AB . In the same way, on the other side of the plane a perpendicular can be drawn parallel to BA produced.

If AB does not meet MN , then at least in one direction it diverges from MN . Through H , any point of the projection of AB on the plane, we can draw a line, HK , parallel to AB towards that part of AB which diverges from MN , and then draw CD parallel to this line and perpendicular to the plane.


Unless AB is parallel to MN it will meet the plane at some point, or the plane and line will have a common perpendicular, and the line will diverge from the plane in both directions.

In the latter case there are two perpendiculars that are parallel to the line, one parallel towards each part of the line.

Two perpendiculars cannot be parallel towards the same part of a line; for then they would be parallel to each other, and two lines cannot be perpendicular to a plane and parallel to each other.

II. BOUNDARY-CURVES AND SURFACES, AND EQUI-DISTANT-CURVES AND SURFACES

Having given the line AB , at its extremity, A , we take any arbitrary angle and produce the side AC so that the perpendicular erected at its middle point shall be parallel to AB . The locus of the point C is a curve which is called oricycle, or boundary-curve. AB is its axis.




From their definition it follows that all boundary-curves are equal, and the boundary-curve is symmetrical with respect to its axis; if revolved through two right angles about its axis, it will coincide with itself.

1. Theorem. *Any line parallel to the axis of a boundary-curve may be taken for axis.*

Let AB be the axis and CD any line parallel to AB , to prove that CD may be taken as axis.


Proof. Draw AC ; also to E , any other point on the curve, draw AE and CE . The perpendiculars erected at the middle points of AC and of AE are parallel to AB and CD and to each



other. Therefore, the perpendicular erected at the middle point of CE , the third side of the triangle ACE , is parallel to them and to CD . CD then may be taken as axis.

Corollary. *The boundary-curve may be slid along on itself without altering its shape; that is, it has a constant curvature.*

2. Theorem. *Two boundary-curves having a common set of axes cut off the same distance on each of the axes, and the ratio of corresponding arcs depends only on this distance.*



Proof. Take any two axes and a third axis bisecting the arc which the first two intercept on one of the two boundary-

curves. By revolving the figure about this axis we show that the curves cut off equal distances on the two axes.

Let AA' , BB' , and CC' be any three axes of the two boundary-curves AB and $A'B'$; let their common length be x and let them intercept arcs s and t on AB , s' and t' on $A'B'$.


When $s = t$, $s' = t'$, and, in general,

$$\frac{s}{t} = \frac{s'}{t'},$$

as we prove, first when s and t are commensurable, and then by the method of limits when they are incommensurable.

The ratio $\frac{s}{s'}$ is, therefore, a constant for the given value of x .

Write $\frac{s}{s'} = f(x)$.



From three boundary-curves having the same set of axes, we find

$$f(x + y) = f(x)f(y).$$

This property is characteristic of the exponential function whose general form is $f(x) = e^{ax}$.* Therefore, $\frac{s}{s'} = e^{ax}$, the value of a depending on the unit of measure (see below p. 76).

* Putting $y = x, 2x, \dots (n-1)x$ in succession, we find

$$f(nx) = [f(x)]^n$$

for positive integer values of n , x being any positive quantity.


Now

$$f\left(\frac{r}{s}x\right) = \left[f\left(\frac{x}{s}\right)\right]^r,$$

and this is the r th power of the s th root of the first member of the equation


$$\left[f\left(\frac{x}{s}\right)\right]^s = f(x); \quad \therefore f\left(\frac{r}{s}x\right) = [f(x)]^{\frac{r}{s}}.$$

3. Theorem. *The area enclosed by two boundary-curves having the same axes and by two of their common axes is proportional to the difference of the intercepted arcs.*



Proof. Let s and s' be the lengths of the intercepted arcs, and l the distance measured on an axis between them. Let t , t' , and k be the corresponding quantities for a second figure constructed in the same way.

If the corresponding lines in the two figures are all equal, the areas are equal, for they can be made to coincide. If only $k = l$, the areas are to each other as corresponding arcs, say as $s' : t'$, proved first when the arcs are commensurable, and then by the method of limits when they are incommensurable.



When l and k are commensurable, suppose

$$\frac{l}{m} = \frac{k}{n} = a.$$

Thus, assuming that $f(x)$ is a continuous function of x , we have proved that for all real positive values of x and n

$$f(nx) = [f(x)]^n,$$

and if we put x for n and 1 for x , we have

$$f(x) = [f(1)]^x.$$

We will write $f(1) = e^a$; then

$$f(x) = e^{ax}.$$

We can draw a series of boundary-curves at distances equal to a on the axes and divide the areas into m and n parts, respectively. If r is the ratio of arcs corresponding to the distance a , these parts will be proportional to the quantities

$$s', s'r, s'r^2, \dots s'r^{m-1};$$

$$t', t'r, t'r^2, \dots t'r^{n-1}.$$

The two areas are then to each other in the ratio

$$s' \frac{r^m - 1}{r - 1} : t' \frac{r^n - 1}{r - 1}.$$

But


$$s'r^m = s \text{ and } t'r^n = t,$$

so that this is the same as the ratio

$$s - s' : t - t'.$$

When l and k are incommensurable, we proceed as in other similar demonstrations.

This theorem is analogous to the one which we have proved about polygons: the area is proportional to the amount of rotation in excess of four right angles in going around the figure, for the rate of rotation in going along a boundary-curve is constant.




The locus of points at a given distance from a straight line is a curve which may be called an equidistant-curve. The perpendiculars from the different points of this curve upon the base line are equal and may be called axes of the curve.

An equidistant-curve fits upon itself when revolved through two right angles about one of its axes or when slid along upon itself. It has a constant curvature.

It can be proved, exactly as in the case of two boundary-curves having the same set of axes, that arcs on an equidistant-curve are proportional to the segments cut off by the axes at their extremities on the base line or on any other equidistant-curve having the same set of axes.

4. Theorem. *The boundary-curve is a limiting curve between the circle and the equidistant-curve; it may be regarded as a circle with infinitely large radius, or as an equidistant-curve whose base line is infinitely distant.*



Proof. Take a line of given length, $AB = 2a$ say, making an angle, A , with a fixed line, AC . Construct another angle at B equal to the angle A , and draw a perpendicular to AB at its middle point, D .

If the angle at A is sufficiently small, we have an isosceles triangle with AB for base, and its vertex at a point, F , on AC . With F as centre, we can draw a circle through the points A and B . Now let the angle at A gradually increase, the rest of the figure varying so as to keep the construction. F will move off indefinitely, and when $A = \Pi(a)$ the three lines AF , BF , and DF will become parallel, and B will become a point on the boundary-curve AB' , which has AC for axis.

On the other hand, if the angle at A were taken acute, but greater than $\Pi(a)$, we should have three lines, AE , BH , and

DF , perpendicular to a line, EH , the base line of an equidistant-curve through the points A and B . Now let the angle A gradually decrease, the rest of the figure varying so as to preserve the construction. The quadrilateral $ADFE$, having three right angles and the fourth angle A decreasing, must increase in area. We get this same movement if we think of AD and DF remaining fixed in the plane while AE revolves about A , making the angle A decrease. Thus the only way in which the area of the quadrilateral can increase is for EH to move off along on AC and become more and more remote from A . When A becomes equal to $\Pi(a)$, BH and DF become parallel to AC , and B falls on the boundary-curve AB' .

Calling the radius of a circle axis, we find that circles, boundary-curves, and equidistant-curves have many properties in common :

The perpendicular erected at the middle point of any chord is an axis. In particular, a tangent is perpendicular to the axis drawn from its point of contact. These are curves cutting at right angles a system of lines through a point, a system of parallel lines, and the perpendiculars to a given line, respectively.


Two of these curves having the same set of axes cut off equal lengths on all these axes, and the ratio of corresponding arcs on two such curves is a constant depending only on the way in which they divide the axes.

Three points determine one of these curves; that is, through any three points not in a straight line we can draw a curve which shall be either a circle, a boundary-curve, or an equidistant-curve, and through any three points only one such curve can be drawn. Any triangle may be inscribed in one and only one of these curves.

Each of these curves can be moved on itself or revolved about any axis through 180° into coincidence with itself.

A boundary-surface or orisphere is a surface generated by the revolution of a boundary-curve about one of its axes.

5. Theorem. *Any line parallel to the axis of a boundary-surface may be regarded as axis.*



Let AA' be the axis, meeting the surface at A , and BB' a line parallel to the axis through any other point, B , of the surface; to prove that BB' may be regarded as axis.

Proof. Let C be a third point on the surface. Draw CC' through C , and through D , E , and F , the middle points of the sides of the plane triangle ABC , draw DD' , EE' , and FF' all parallel to AA' . Finally, let OO' be parallel to these lines and perpendicular to the plane ABC . The projecting planes of the other parallels all pass through OO' (see I, 9).

Since AA' is axis to the surface, EE' and FF' are perpendicular to AC and AB , respectively. Draw FK perpendicular to the plane ABC at F . It will lie in the projecting plane OFF' . AB , being perpendicular to FF' and to FK , is perpen-

dicular to this plane, OFF' , and therefore to OF . In the same way we prove that AC is perpendicular to OE . Therefore, BC is perpendicular to OD (Chap. I, I, 5). But OD is the intersection of the plane ABC with the plane ODD' . Hence, BC is perpendicular to this plane and to DD' (Chap. I, II, 15).

DD' being parallel to BB' lies in the plane determined by BB' and BC , and in this plane only one perpendicular can be drawn to BC at its middle point. Therefore, if we pass any plane through BB' and from B draw a chord to any other point, C , of its intersection with the surface, the perpendicular in this plane to BC , erected at the middle point of BC , will be parallel to BB' . This proves that the section is a boundary-curve, having BB' for axis, and that the surface can be generated by the revolution of such a boundary-curve around BB' .

Therefore, BB' may be regarded as axis of the surface.

A plane passed through an axis of a boundary-surface is called a principal plane. Every principal plane cuts the surface in a boundary-curve. Any other plane cuts the surface in a circle; for the surface may be regarded as a surface of revolution having for axis of revolution that axis which is perpendicular to the plane. This perpendicular may be called the axis of the circle, and the point where it meets the surface, the pole of the circle. The pole of a circle on a boundary-surface is at the same distance from all the points of the circle, distance being measured along boundary-lines on the surface.

Any two boundary-surfaces can be made to coincide, and a boundary-surface can be moved upon itself, any point to the position of any other point, and any boundary-curve through the first point to the position of any boundary-curve through the second point. We may say that a boundary-surface has a constant curvature, the same for all these surfaces. Figures on a boundary-surface can be moved about or put upon any other boundary-surface without altering their shape or size.

We can develop a Geometry on the boundary-surface. By *line* we mean the boundary-curve in which the surface is cut by a principal plane. The angle between two lines is the same as the dihedral angle between the two principal planes which cut out the lines on the surface.

6. Theorem. *Geometry on the boundary-surface is the same as the ordinary Euclidean Plane Geometry.*


Proof. On two boundary-surfaces with the same system of parallel lines for axes corresponding triangles are similar; that is, corresponding angles are equal, having the same measures as the dihedral angles which cut them out, and corresponding lines are proportional by (2). But we can place these figures on the same surface; therefore, on one boundary-surface we can have similar triangles. Thus, we can diminish the sides of a triangle without altering their ratios or the angles. We can do this indefinitely; for the ratio of corresponding lines on the two surfaces, being expressed by the function e^{ax} of the distance between them, can be made as large as we please by taking x sufficiently large. If we assume that figures on the boundary-surface become more and more like plane figures when we diminish indefinitely their size, it follows that a triangle on this surface approaches more and more the form of an infinitesimal plane triangle, for which the sum of the angles is two right angles, and the angles and sides have the same relations as in the Euclidean Plane Geometry. All the formulæ of Plane Trigonometry with which we are familiar hold, then, for triangles on the boundary-surface.

On the boundary-surface we have the "hypothesis of the right angle." Rectangles can be formed, and the area of a rectangle is proportional to the product of its base and altitude, while the area of a triangle is half of the area of a rectangle having the same base and altitude.

An equidistant-surface is a surface generated by the revolution of an equidistant-curve about one of its axes. It is the locus of points at a given perpendicular distance from a plane. Any perpendicular to the plane may be regarded as an axis, and the surface is a surface cutting at right angles a system of lines perpendicular to the plane. The surface has a constant curvature, fitting upon itself in any position.

III. TRIGONOMETRICAL FORMULÆ

1. Let ABC be a plane right triangle. Erect AA' perpendicular to its plane and draw BB' and CC' parallel to AA' . Draw a boundary-surface through A , having these lines for axes and forming the boundary-surface triangle $AB''C''$. Also construct the spherical triangle about the point B .



The angle A is the same in the plane triangle and in the boundary-surface triangle. The planes through AA' are perpendicular to ABC . Hence, the spherical triangle has a right angle at the vertex which lies on c , and BC being perpendicular to CA is perpendicular to the plane of CC' and AA' . Therefore, the plane BCC' is perpendicular to the plane ACC' ,

and the dihedral whose edge is BC has for plane angle the angle $ACC' = \Pi(b)$. Since the boundary-surface triangle is right-angled at C'' , the angle B'' , or what is the same thing, the dihedral whose edge is BB' , is the complement of the angle A .

In the spherical triangle the side opposite the right angle is $\Pi(a)$, the two sides about the right angle are $\Pi(c)$ and B , and the opposite angles are $\Pi(b)$ and $90^\circ - A$.

Applying to these quantities the trigonometrical formulæ for spherical right triangles, we get at once the relations that connect the sides and angles of plane right triangles.

Produce to quadrants the two sides about the angle whose value is the complement of A . We form in this way a spherical right triangle in which the side opposite the right angle is the complement of $\Pi(c)$, the two sides about the right angle are the complements of $\Pi(a)$ and $\Pi(b)$, and their opposite angles are the complements of B and A . From this triangle we deduce the following rule for passing from the formulæ of spherical right triangles to those of plane triangles :

Interchange the two angles (or the two sides) and everywhere use the complementary function, taking the corresponding angle of parallelism for the sides.

The formulæ for spherical right triangles are

$$\begin{array}{ll} \sin A = \frac{\sin a}{\sin c} & \sin B = \frac{\sin b}{\sin c} \\ \cos A = \frac{\tan b}{\tan c} & \cos B = \frac{\tan a}{\tan c} \\ \tan A = \frac{\tan a}{\sin b} & \tan B = \frac{\tan b}{\sin a} \\ \sin A = \frac{\cos B}{\cos b} & \sin B = \frac{\cos A}{\cos a} \end{array}$$

$$\cos c = \cos a \cos b.$$

$$\cos c = \cot A \cot B.$$

From these, by the rule given on the previous page, we derive the following formulæ for plane right triangles:

$$\cos B = \frac{\cos \Pi(a)}{\cos \Pi(c)} \quad \cos A = \frac{\cos \Pi(b)}{\cos \Pi(c)}.$$

$$\sin B = \frac{\cot \Pi(b)}{\cot \Pi(c)} \quad \sin A = \frac{\cot \Pi(a)}{\cot \Pi(c)}.$$

$$\cot B = \frac{\cot \Pi(a)}{\cos \Pi(b)} \quad \cot A = \frac{\cot \Pi(b)}{\cos \Pi(a)}.$$

$$\cos B = \frac{\sin A}{\sin \Pi(b)} \quad \cos A = \frac{\sin B}{\sin \Pi(a)}.$$

$$\sin \Pi(c) = \sin \Pi(a) \sin \Pi(b).$$

$$\sin \Pi(c) = \tan A \tan B.*$$


We can obtain the formulæ for oblique plane triangles by dropping a perpendicular from one vertex upon the opposite side, thus forming two right triangles.

2. Take the relation


$$\sin \Pi(a) = \frac{\sin B}{\cos A}.$$

Let p , q , and r be the sides of the triangle $AB''C''$ of our last demonstration and p' , q' , and r the corresponding sides

* We can arrange the parts of a right triangle so as to apply Napier's rules; namely, the arrangement would be




of the triangle formed in the same way on a boundary-surface tangent to the plane ABC at B .



$$\sin B = \frac{q'}{r},$$

$$\cos A = \frac{q}{r}.$$

$$\therefore \sin \Pi(a) = \frac{q'}{q}.$$




Now q and q' are corresponding arcs on two boundary-curves which have the same set of parallel lines as axes, and their distance apart, x , is the distance from a boundary-curve of the extremity of a tangent of arbitrary length, a . Thus, we have for corresponding arcs

$$\frac{s'}{s} = \sin \Pi(a).$$

3. To MN , a given straight line, erect a perpendicular at a point, O , and on this perpendicular lay off $OA = y$ below MN , and OB and BP each equal to x above MN , x and y being any arbitrary lengths. At P draw PR perpendicular to OP and

extending towards the left, and through B draw EF making with OP an angle $\Pi(x)$, and therefore parallel on one side to ON and on the other side to PR . Finally, draw AK and AH , the two parallels to EF through A .



At the point A we have four angles of parallelism :

$$CAK = CAH = \Pi(AC),$$

$$OAK = \Pi(y),$$

$$PAH = \Pi(y + 2x).$$

Therefore, $\Pi(y) = \Pi(AC) + BAC,$

and $\Pi(y + 2x) = \Pi(AC) - BAC.$

Now in the right triangle ABC

$$\cos \Pi(y + x) = \frac{\cos \Pi(AC)}{\cos BAC},$$

or
$$\frac{1 - \cos \Pi(y + x)}{1 + \cos \Pi(y + x)} = \frac{\cos BAC - \cos \Pi(AC)}{\cos BAC + \cos \Pi(AC)}$$

$$= \frac{\sin \frac{1}{2}[\Pi(AC) + BAC] \sin \frac{1}{2}[\Pi(AC) - BAC]}{\cos \frac{1}{2}[\Pi(AC) + BAC] \cos \frac{1}{2}[\Pi(AC) - BAC]};$$

whence,

$$\tan^2 \frac{1}{2} \Pi(y+x) = \tan \frac{1}{2} \Pi(y) \tan \frac{1}{2} \Pi(y+2x).$$

$\tan \frac{1}{2} \Pi(x)$ is then a function of x , say $f(x)$, satisfying the condition

$$[f(y+x)]^2 = f(y)f(y+2x),$$

or

$$\frac{f(y+x)}{f(y)} = \frac{f(y+2x)}{f(y+x)},$$

and putting successively in this equation $y+x$, $y+2x$, etc., for y , we may add

$$= \frac{f(y+3x)}{f(y+2x)} = \dots = \frac{f(y+nx)}{f[y+(n-1)x]}.$$

$\Pi(0) = \frac{\pi}{2}$ and $\tan \frac{1}{2} \Pi(0) = 1$; therefore, putting $y = 0$ in the first and last of all these fractions, we have

$$f(x) = \frac{f(nx)}{f[(n-1)x]},$$

or

$$f(nx) = f[(n-1)x]f(x).$$

$$\therefore f(nx) = [f(x)]^n.$$

This equation is characteristic of the exponential function.* $\Pi(x)$ being an acute angle, $\tan \frac{1}{2} \Pi(x) < 1$; therefore, we may write $f(1) = e^{-a'}$, so that $f(x) = e^{-a'x}$. a' depends on the unit of measure; we will take the unit so that $a' = 1$. Finally, since $\Pi(-x) = \pi - \Pi(x)$,

$$\tan \frac{1}{2} \Pi(-x) = \cot \frac{1}{2} \Pi(x) = [\tan \frac{1}{2} \Pi(x)]^{-1}.$$

That is, for all real values of x

$$\tan \frac{1}{2} \Pi(x) = e^{-x},$$

* See footnote, p. 45.

or
$$\frac{1 - \cos \Pi(x)}{\sin \Pi(x)} = \cos ix + i \sin ix.*$$

* i stands for $\sqrt{-1}$. The best way to get the relations between the exponential and trigonometrical functions is by their developments in series :

$$\begin{aligned} e^x &= 1 + x + \frac{x^2}{2} + \dots + \frac{x^n}{n} + \dots, \\ \cos x &= 1 - \frac{x^2}{2} + \frac{x^4}{4} - \dots + (-1)^n \frac{x^{2n}}{2n} + \dots, \\ \sin x &= x - \frac{x^3}{3} + \frac{x^5}{5} - \dots + (-1)^n \frac{x^{2n+1}}{2n+1} + \dots. \end{aligned}$$

These series are convergent for all values of x .

Putting ix for x , we have

$$\begin{aligned} e^{ix} &= 1 + ix - \frac{x^2}{2} - \frac{ix^3}{3} + \dots \\ &= 1 - \frac{x^2}{2} + \frac{x^4}{4} - \dots + i \left(x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \right); \end{aligned}$$

i.e.,

$$e^{ix} = \cos x + i \sin x.$$

Also

$$e^{-ix} = \cos x - i \sin x.$$

$$\therefore \cos x = \frac{1}{2}(e^{ix} + e^{-ix}),$$

$$\sin x = \frac{1}{2i}(e^{ix} - e^{-ix}).$$

Again, putting ix for x , we have

$$e^x = \cos ix - i \sin ix,$$

$$e^{-x} = \cos ix + i \sin ix;$$

and

$$\cos ix = \frac{1}{2}(e^x + e^{-x}),$$

$$\sin ix = -\frac{1}{2i}(e^x - e^{-x}).$$

$$\cos ix = 1 + \frac{x^2}{2} + \frac{x^4}{4} + \dots,$$

$$\sin ix = ix \left(1 + \frac{x^2}{3} + \frac{x^4}{5} + \dots \right).$$

For real values of x , $\cos ix$ and $\frac{\sin ix}{ix}$ are real and positive, and vary from 1 to ∞ as x varies from 0 to ∞ .

In the equation $\cos^2 ix + \sin^2 ix = 1$, the first term is real and positive for real values of x , the second term is real and negative; therefore, $\sin ix$ is in absolute value less than $\cos ix$, and $\tan ix$ is in absolute value less than 1. $\tan ix$ varies in absolute value from 0 to 1 as x varies from 0 to ∞ .

Changing the sign of x , we have

$$\frac{1 + \cos \Pi(x)}{\sin \Pi(x)} = \cos ix - i \sin ix,$$

and, adding and subtracting,

$$\begin{aligned}\frac{1}{\sin \Pi(x)} &= \cos ix, \\ \cot \Pi(x) &= -i \sin ix.\end{aligned}$$

The nature of the angle of parallelism is, therefore, expressed by the equations

$$\begin{aligned}\sin \Pi(x) &= \frac{1}{\cos ix}, \\ \tan \Pi(x) &= \frac{i}{\sin ix}, \\ \cos \Pi(x) &= \frac{\tan ix}{i}.\end{aligned}$$

4. Substituting in the formulæ of plane right triangles, we find that they reduce to those of spherical right triangles with ia , ib , and ic for a , b , and c , respectively. The formulæ of oblique triangles are obtained from those of right triangles in the same way as on the sphere, and thus all the formulæ of Plane Trigonometry are obtained from those of Spherical Trigonometry simply by making this change.

As fundamental formulæ for oblique triangles we write

$$\begin{aligned}\frac{\sin A}{\sin ia} &= \frac{\sin B}{\sin ib} = \frac{\sin C}{\sin ic}, \\ \cos ia &= \cos ib \cos ic + \sin ib \sin ic \cos A, \\ \cos A &= -\cos B \cos C + \sin B \sin C \cos ia.\end{aligned}$$

In the notation of the Π -function, these are

$$\sin A \tan \Pi(a) = \sin B \tan \Pi(b) = \sin C \tan \Pi(c),$$

$$\frac{\sin \Pi(b) \sin \Pi(c)}{\sin \Pi(a)} = 1 - \cos \Pi(b) \cos \Pi(c) \cos A,$$

$$\cos A = -\cos B \cos C + \frac{\sin B \sin C}{\sin \Pi(a)}.$$

5. Since for very small values of x we have approximately

$$\sin ix = ix,$$

$$\cos ix = 1 + \frac{x^2}{2},$$

$$\tan ix = ix,$$

our formulæ for infinitesimal triangles reduce to

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c},$$

$$a^2 = b^2 + c^2 - 2bc \cos A,$$


$$\cos A = -\cos(B + C).$$

6. Triangles on an equidistant-surface are similar to their projections on the base plane; that is, they have the same angles and their sides are proportional. Thus the formulæ of Plane Trigonometry hold for any equidistant-surface if with the letters representing the sides we put, besides i , a constant factor depending on the distance of the surface from the plane.

CHAPTER III

THE ELLIPTIC GEOMETRY

In the hypothesis of the obtuse angle a straight line is of finite length and returns into itself. This length is the same for all lines, since any two lines can be made to coincide. Two straight lines always intersect, and two lines perpendicular to a third intersect at a point whose distance from the third on either line is half the entire length of a straight line.



1. A straight line does not divide the plane. Starting from the point of intersection of two lines and passing along one of them a certain finite distance, we come to the intersection point again without having crossed the other line. Thus, we can pass from one side of the line to the other without having crossed it.

There is one point through which pass all the perpendiculars to a given line. It is called the pole of that line, and the line is its polar. Its distance from the line is half the entire length of a straight line, and the line is the locus of points at this distance from its pole. Therefore, if the pole of one

line lies on another, the pole of the second lies on the first, and the intersection of two lines is the pole of the line joining their poles.

The locus of points at a given distance from a given line is a circle having its centre at the pole of the line. The straight line is a limiting form of a circle when the radius becomes equal to half the entire length of a line.

We can draw three lines, each perpendicular to the other two, forming a trirectangular triangle. It is also a self-polar triangle; each vertex is the pole of the opposite side.

2. All the perpendiculars to a plane in space meet at a point which is the pole of the plane. It is the centre of a system of spheres of which the plane is a limiting form when the radius becomes equal to half the entire length of a straight line.

Figures on a plane can be projected from similar figures on any sphere which has the pole of the plane for centre. That is, they have equal angles and corresponding sides in a constant ratio that depends only on the radius of the sphere. Two corresponding angles are equal, because they are the same as the dihedral angles formed by the two planes through the centre of the sphere which cut the sphere and the plane in the sides of the angles. Corresponding lines are proportional; for if two arcs on the sphere are equal, their projections on the plane are equal; and that, in general, two arcs have the same ratio as their projections on the plane is proved, first when they are commensurable, and by the method of limits when they are incommensurable.

Geometry on a plane is, therefore, like Spherical Geometry, but the plane corresponds to only half a sphere, just as the diameters of a sphere correspond to the points of half the surface. Indeed, the points and straight lines of a plane correspond exactly to the lines and planes through a point,

but we can realize the correspondence better that compares the plane with the surface of a sphere. If we can imagine that the points on the boundary of a hemisphere at opposite extremities of diameters are coincident, the hemisphere will correspond to the elliptic plane. There is no particular line of the plane that plays the part of boundary. All lines of the plane are alike; the plane is unbounded, but not infinite in extent.


The entire straight line corresponds to a semicircle. We will take such a unit for measuring length that the entire length of a line shall be π ; the formulæ of Spherical Trigonometry will then apply without change to our plane. Distances on a line will then have the same measure as the angles which they subtend at the pole of the line, and the angle between two lines will be equal to the distance between their poles. The distance from any point to its polar, half the entire length of a straight line, may then be called a quadrant.

We can form a self-polar tetraëdron by taking three mutually perpendicular planes and the plane which has their intersection for pole. The vertices of this tetraëdron are the poles of the opposite faces. At each vertex is a trirectangular trihedral, and each face is a trirectangular triangle.

3. Theorem. *All the planes perpendicular to a fixed line intersect in another fixed line, called its polar or conjugate. The relation is reciprocal, and all the points of either line are at a quadrant's distance from all the points of the other.*

Proof. Let the two planes perpendicular to the line AB at H and K intersect in CD . Pass a plane through AB and R , any point of CD . This plane will intersect the two given planes in HR and KR . HR and KR are perpendicular to AB ; therefore, R is at a quadrant's distance from H and K . R is then the pole of AB in the plane determined by AB and R ,

and is at a quadrant's distance from every point of AB . But R is any point of CD ; therefore, any point of either line is at a quadrant's distance from each point of the other line, and a point which is at a quadrant's distance from one line lies in



the other line. Again, any point, H , of AB , being at a quadrant's distance from all the points of CD , is the pole of CD in the plane determined by it and CD . Thus, HR and KR are both perpendicular to CD , and the plane determined by AB and R is perpendicular to CD .


The opposite edges of a self-polar tetraëdron are polar lines.

All the lines which intersect a given line at right angles intersect its polar at right angles. Therefore, the distances of any point from two polar lines are measured on the same straight line and are together equal to a quadrant. Two points which are equidistant from one line are equidistant from its polar.

The locus of points which are at a given distance from a fixed line is a surface of revolution having both this line and its polar as axes. We may call it a surface of double revolution. The parallel circles about one axis are meridian curves for the other axis. If a solid body, or, we may say, all space, move along a straight line without rotating about it, it will rotate about the conjugate line as an axis without sliding

along it. A motion along a straight line combined with a rotation about it is called a screw motion. A screw motion may then be described as a rotation about each of two conjugate lines or as a sliding along each of two conjugate lines.

4. Theorem. *In the elliptic geometry there are lines not in the same plane which have an infinite number of common perpendiculars and are everywhere equidistant.*



Given any two lines in the same plane and their common perpendicular. If we go out on these lines in either direction from the perpendicular, they approach each other. Now revolve one of them about this perpendicular so that they are no longer in the same plane. After a certain amount of rotation the lines will have an infinite number of common perpendiculars and be equidistant throughout their entire length.

Proof. Let p be the length of the common perpendicular AC , and take points B and D on the two lines on the same side of this perpendicular at a distance, a .

$BD < p$, but if CD revolve about AC , BD will become longer than p by the time CD is revolved through a right angle; for BCD will then be a right triangle, with BD for hypotenuse and BC , the hypotenuse of the triangle ABC , for one of its sides, so that we shall have $BD > BC$ and $BC > AC$.

Suppose, when CD has revolved through an angle, θ , BD becomes equal to p and takes the position BD' . The triangles ABC and $D'BC$ are equal, having corresponding sides equal. Therefore, BD' is perpendicular to CD' . BD' is also perpendicular to BA ; for if we take the diedral $A-BC-D'$ and place it upon itself so that the positions of B and C shall be interchanged, A will fall on the position of D' , and D' on the position of A , and the angle $D'BA$ must equal the angle ACD' . Therefore, BD' as well as CA is a common perpendicular to the lines AB and CD' .

Now at the point C we have a triedral whose three edges are CB , CD , and CD' . Moreover, the diedral along the edge CD is a right diedral; therefore, the three face angles of the triedral satisfy the same relations as do the three sides of a spherical right triangle; namely,

$$\cos BCD' = \cos BCD \cos DCD'.$$

$$\text{But} \quad BCD = \frac{\pi}{2} - ACB \quad \text{and} \quad BCD' = ABC.$$

Hence, this relation may be written

$$\cos ABC = \sin AC'B \cos \theta.$$

Again, in the right triangle ABC'

$$\sin AC'B = \frac{\cos ABC}{\cos p}.$$

$$\therefore \cos \theta = \cos p,$$

or, since θ and p are less than $\frac{\pi}{2}$,

$$\theta = p.$$

The angle θ , therefore, does not depend upon a . If we take any two lines in a plane and turn one about their common perpendicular through an angle equal in measure to the length

of that perpendicular, the two lines will then be everywhere equidistant.

As we have no parallel lines in the ordinary sense in this Geometry, the name *parallel* has been applied to lines of this kind. They have many properties of the parallel lines of Euclidean Geometry.


Through any point two lines can be drawn parallel to a given line. These are of two kinds, sometimes distinguished as right-wound and left-wound. They lie entirely on a surface of double revolution, having the given line as axis. The surface is, therefore, a ruled surface and has on it two sets of rectilinear generators like the hyperboloid of one sheet.

CHAPTER IV

ANALYTIC NON-EUCLIDEAN GEOMETRY

WE shall use the ordinary polar coördinates, ρ and θ , and for the rectangular coördinates, x and y , of a point, we shall use the intercepts on the axes made by perpendiculars through the point to the axes. The formulæ depend upon the trigonometrical relations, and in our two Geometries differ only in the use of the imaginary factor i with lengths of lines.

I. HYPERBOLIC ANALYTIC GEOMETRY



1. The relations between polar and rectangular coördinates :

The angles at the origin which the radius vector makes with the axes are complementary. From the two right triangles we have


$$\tan ix = \cos \theta \tan i\rho,$$

$$\tan iy = \sin \theta \tan i\rho.$$

Therefore,

$$\tan^2 i\rho = \tan^2 ix + \tan^2 iy,$$


$$\tan \theta = \frac{\tan iy}{\tan ix}.$$



2. The distance, δ , between two points :

$$\cos i\delta = \cos i\rho \cos i\rho' + \sin i\rho \sin i\rho' \cos (\theta' - \theta).$$

δ and one of the points being fixed, this may be regarded as the polar equation of a circle.



3. The equation of a line :

Let p be the length of the perpendicular from the origin upon the line, and α the angle which the perpendicular makes

with the axis of x . From the right triangle formed with this perpendicular and ρ we have

$$\tan ip \cos (\theta - \alpha) = \tan ip.$$

This is the polar equation of the line. We get the equation in x and y by expanding and substituting; namely,


$$\cos \alpha \tan ix + \sin \alpha \tan iy = \tan ip.$$

The equation $a \tan ix + b \tan iy = i$ represents a line for which

$$a^2 + b^2 = \frac{-1}{\tan^2 ip}.$$

Now, for real values of p , $-\tan^2 ip < 1$ (see footnote, p. 59). The line is therefore real if a and b are real, and if

$$a^2 + b^2 > 1.$$



4. The distance, δ , of a point from a line:

Let the radius vector to the point intersect the line at A , and let ρ_1 be the radius vector to A . We have two right

triangles with equal angles at A , and from the expressions for the sines of these angles we get the equation

$$\frac{\sin i\delta}{\sin i(\rho - \rho_1)} = \frac{\sin ip}{\sin i\rho_1}.$$

This equation holds for all points, $x y$, of the plane, δ being negative when the point is on the same side of the line as the origin, and zero when the point is on the line.


$$\sin i\delta = \frac{\sin ip}{\tan i\rho_1} \sin i\rho - \sin ip \cos i\rho.$$

Now,
$$\tan i\rho_1 = \frac{\tan ip}{\cos(\theta - \alpha)}.$$

$$\therefore \frac{\sin ip}{\tan i\rho_1} \sin i\rho = \sin i\rho \cos i\rho \cos(\theta - \alpha),$$

and
$$\sin i\delta = \cos i\rho \cos ip [\tan i\rho \cos(\theta - \alpha) - \tan ip].$$

δ being fixed, this may be regarded as the polar equation of an equidistant-curve.



5. The angle between two lines :

ϕ being the angle which a line makes with the radius vector at any point, we have

$$\cos \phi = \cos ip \sin(\theta - \alpha),$$

$$\sin \phi = \frac{\sin ip}{\sin i\rho}.$$

For two lines intersecting at this point,

$$\begin{aligned}\sin \phi_1 \sin \phi_2 &= \frac{\sin ip_1 \sin ip_2}{\sin^2 i\rho} \\ &= \sin ip_1 \sin ip_2 + \frac{\sin ip_1 \sin ip_2}{\tan^2 i\rho}.\end{aligned}$$

Now, from the equation of the line

$$\frac{\sin ip_1}{\tan i\rho} = \cos ip_1 \cos(\theta - \alpha_1),$$

$$\frac{\sin ip_2}{\tan i\rho} = \cos ip_2 \cos(\theta - \alpha_2);$$

$$\begin{aligned}\text{so that} \quad \sin \phi_1 \sin \phi_2 &= \sin ip_1 \sin ip_2 \\ &\quad + \cos ip_1 \cos ip_2 \cos(\theta - \alpha_1) \cos(\theta - \alpha_2).\end{aligned}$$

$$\text{Again,} \quad \cos \phi_1 \cos \phi_2 = \cos ip_1 \cos ip_2 \sin(\theta - \alpha_1) \sin(\theta - \alpha_2).$$

Adding these equations, we have

$$\cos(\phi_2 - \phi_1) = \sin ip_1 \sin ip_2 + \cos ip_1 \cos ip_2 \cos(\alpha_2 - \alpha_1).$$

Two lines are perpendicular if

$$\cos(\alpha_2 - \alpha_1) + \tan ip_1 \tan ip_2 = 0.$$

$$\text{The lines} \quad a \tan ix + b \tan iy = i,$$

$$a' \tan ix + b' \tan iy = i$$

$$\text{are perpendicular if} \quad aa' + bb' = 1.$$

6. The equation of a circle in x and y :


$$\sin i\rho \cos \theta = \cos i\rho \tan ix,$$

$$\sin i\rho \sin \theta = \cos i\rho \tan iy;$$

$$\text{also,} \quad \cos i\rho = \frac{1}{\sqrt{1 + \tan^2 i\rho}} = \frac{1}{\sqrt{1 + \tan^2 ix + \tan^2 iy}}.$$

The equation of a circle may, therefore, be written

$$(1 + \tan^2 ix + \tan^2 iy)(1 + \tan^2 ix' + \tan^2 iy') \cos^2 i\delta \\ = (1 + \tan ix \tan ix' + \tan iy \tan iy')^2.$$



7. The equation of a boundary-curve :

Let the axis of the boundary-curve which passes through the origin make an angle, α , with the axis of x , and let the point where the boundary-curve cuts this axis be at a distance, k , from the origin, positive if the origin is on the convex side of the curve, negative if the origin is on the concave side of the curve. The boundary-curve is the limiting position of a circle whose centre, on this axis, moves off indefinitely.

ρ' being the radius vector to the centre, the radius of the circle is $\rho' - k$, and its equation may be written

$$\cos i(\rho' - k) = \cos i\rho \cos i\rho' + \sin i\rho \sin i\rho' \cos(\theta - \alpha),$$

or, expanding and dividing by $\cos i\rho'$,

$$\cos ik + \tan i\rho' \sin ik = \cos i\rho + \sin i\rho \tan i\rho' \cos(\theta - \alpha).$$

Now, let ρ' increase indefinitely. $\tan i\rho'$ tends to the limit i , so that the limit of the first member of the equation is


$$\cos ik + i \sin ik, \text{ or } e^{-k},$$

and the polar equation of the curve is

$$e^{-k} = \cos i\rho [1 + i \tan i\rho \cos(\theta - \alpha)];$$

or, in xy coördinates,

$$\begin{aligned} (1 + \tan^2 ix + \tan^2 iy) e^{-2k} \\ = (1 + i \cos \alpha \tan ix + i \sin \alpha \tan iy)^2. \end{aligned}$$



Let k be negative and equal, say, to $-b$, and let $\alpha = 0$; also, let a be the ordinate of the point A where the curve cuts the axis of y .

Substituting in the equation, we find

$$e^b = \cos ia.$$

Through A draw a line parallel to the axis of x , and, therefore, making an angle, $\Pi(a)$, with the axis of y . If we draw a boundary-curve through the origin having the same set of parallel lines for axes, so that the two boundary-curves cut

off a distance, b , on these axes, we know that the ratio of corresponding arcs is

$$\frac{s'}{s} = \sin \Pi(\alpha) = \frac{1}{\cos i\alpha}; \quad (\text{See p. 56.})$$

therefore,
$$\frac{s'}{s} = e^{-b}. \quad (\text{See p. 45.})$$

8. The equation of an equidistant-curve :

The polar equation of (4) reduced to an equation in x and y takes the form


$$\begin{aligned} (1 + \tan^2 ix + \tan^2 iy) \sin^2 i\delta \\ = \cos^2 ip (\cos \alpha \tan ix + \sin \alpha \tan iy - \tan ip)^2. \end{aligned}$$

9. Comparison of the three equations :

The equation

$$(1 + \tan^2 ix + \tan^2 iy) c^2 = -(i - a \tan ix - b \tan iy)^2$$

represents a circle, a boundary-curve, or an equidistant-curve, according as $a^2 + b^2 < 1, = 1, > 1$, respectively.



10. Differential formulæ :

Suppose we have an isosceles triangle in which the angle A at the vertex diminishes indefinitely. In the formula


$$\frac{\sin A}{\sin ia} = \frac{\sin C}{\sin ic}$$

we may put for $\sin A, \sin ia, \sin C;$
 $A, ia, 1,$

respectively. Therefore,

$$(I.) \quad ia = \sin ic \cdot A.$$

Corollary. *In a circle of radius r , the ratio of any arc to the angle subtended at the centre is $\sin ir$.*



Again, in the right triangle ABC , let the hypotenuse c revolve about the vertex A . Differentiating the equation

$$\sin A = \frac{\cos B}{\cos ib},$$

where b is constant, we have

$$\cos A dA = - \frac{\sin B dB}{\cos ib}.$$

But

$$\sin B = \frac{\cos A}{\cos ia};$$

$$\therefore dB = - \cos ia \cos ib dA,$$

or (II.)


$$dB = - \cos ic dA.$$

Now, using polar coördinates, we have an infinitesimal right triangle whose hypotenuse, ds , makes an angle, say ϕ , with the radius vector (see figure on page 78). The two sides about the right angle are $d\rho$ and $\frac{\sin i\rho}{i} d\theta$;

therefore,

$$ds^2 = d\rho^2 - \sin^2 i\rho d\theta^2,$$

$$\tan \phi = \frac{\sin i\rho}{i} \frac{d\theta}{d\rho}.$$



For two arcs cutting at right angles, let d' denote differentiation along the second arc :

$$\frac{\sin i\rho}{i} \frac{d\theta}{d\rho} = - \frac{i}{\sin i\rho} \frac{d'\rho}{d'\theta},$$

or

$$\frac{d\rho}{d\theta} \frac{d'\rho}{d'\theta} = \sin^2 i\rho.$$

11. Area :

It equals

$$\iint \frac{\sin i\rho}{i} d\rho d\theta.*$$

We will consider only the case where the origin is within the area to be computed and where each radius vector meets the bounding curve once, and only once.

Integrating with respect to ρ , from $\rho = 0$, we have


$$\int_0^{2\pi} (\cos i\rho - 1) d\theta,$$

or

$$\int_0^{2\pi} \cos i\rho d\theta - 2\pi.$$

* The unit of area being so chosen that the area of an infinitesimal rectangle may be expressed as the product of its base and altitude.

Suppose P and P' are two "consecutive" points on the curve, PM and $P'M'$ the tangents at these points, and ϕ the



angle which the tangent makes with the radius vector. The angle $MP'M'$ indicates the amount of turning or rotation at these points as we go around the curve.

Now, by (II.),

$$MP'M' = d\phi + \cos i\rho d\theta.$$

In going around the curve, ϕ may vary but finally returns to its original value. That is, for our curve

$$\int d\phi = 0,$$

and the amount of rotation is

$$\int_0^{2\pi} \cos i\rho d\theta.$$

Hence, the area is equal to the excess over four right angles in the amount of rotation as we go around the curve. This theorem can be extended to any finite area.

12. A modified system of coördinates :

Our equations take simple forms if we write iu for $\tan ix$, iv for $\tan iy$, ir for $\tan i\rho$, and so on for all lengths of lines.

Thus, we have $u^2 + v^2 = r^2$.*

The equation of a line is

$$au + bv = 1,$$

and the equation


$$(1 - u^2 - v^2)c^2 = (1 - au - bv)^2$$

represents a circle, a boundary-curve, or an equidistant-curve, according as $a^2 + b^2 < 1, = 1, > 1$, respectively.

II. ELLIPTIC ANALYTIC GEOMETRY


The Elliptic Analytic Geometry may be developed just as we have developed the Hyperbolic Analytic Geometry, and the formulæ are the same with the omission of the factor i . But these formulæ are also very easily obtained from the relation of line and pole, and we shall produce them in this way.

The formulæ of Elliptic Plane Analytic Geometry may be applied to a sphere in any of our three Geometries.



1. The relations between polar and rectangular coördinates :

$$\tan x = \cos \theta \tan \rho, \quad \tan y = \sin \theta \tan \rho;$$




* If we draw a quadrilateral with three right angles and the diagonal to the acute angle, and use a, b , and c in the same way that u, v , and r are used above, the five parts lettered in the figure have the relations of a

right triangle in the Euclidean Geometry; e.g.,
 $a^2 + b^2 = c^2, \quad \sin A = \frac{a}{c}, \quad \text{etc.}$

therefore,

$$\tan^2 \rho = \tan^2 x + \tan^2 y,$$

$$\tan \theta = \frac{\tan y}{\tan x}.*$$



2. The distance, δ , between two points :

$$\cos \delta = \cos \rho \cos \rho' + \sin \rho \sin \rho' \cos (\theta' - \theta).$$

This may be regarded as the polar equation of a circle of radius δ , ρ' and θ' being the polar coördinates of the centre.

Now,

$$\sin \rho \cos \theta = \cos \rho \tan x,$$

$$\sin \rho \sin \theta = \cos \rho \tan y;$$


$$\text{also, } \cos \rho = \frac{1}{\sqrt{1 + \tan^2 \rho}} = \frac{1}{\sqrt{1 + \tan^2 x + \tan^2 y}}.$$

The equation of a circle in rectangular coördinates may, therefore, be written

$$(1 + \tan^2 x + \tan^2 y) (1 + \tan^2 x' + \tan^2 y') \cos^2 \delta = (1 + \tan x \tan x' + \tan y \tan y')^2.$$

* The line which has the origin for pole forms with the coördinate axes a trirectangular triangle, and x , y , and θ may be regarded as representing the directions of the given point from its three vertices.

On the sphere, if we take as origin the pole of the equator, ρ and θ are colatitude and longitude. x and y , one with its sign changed, are the bearings of the point from two points 90° apart on the equator.



3. The equation of a line :

When $\delta = \frac{\pi}{2}$, the circle becomes a straight line. For this we have, therefore, the equation

$$\tan x \tan x' + \tan y \tan y' + 1 = 0.$$

$x'y'$ is the pole of the line.

From the equation

$$\tan \rho \cos(\theta - \alpha) = \tan p,$$

or

$$\cos \alpha \tan x + \sin \alpha \tan y = \tan p,$$

we find

$$\tan x' = -\frac{\cos \alpha}{\tan p},$$

$$\tan y' = -\frac{\sin \alpha}{\tan p},$$

as can be shown geometrically, the polar coördinates of this point being

$$p + \frac{\pi}{2}, \alpha.$$

The equation $a \tan x + b \tan y + 1 = 0$ represents a real line for any real values of a and b .

4. The distance, δ , of a point from a straight line :

This is the complement of the distance between the point and the pole of the line; it is expressed by the equation

$$\begin{aligned}\sin \delta &= -\cos \rho \sin p + \sin \rho \cos p \cos (\theta - \alpha) \\ &= \cos \rho \cos p [\tan \rho \cos (\theta - \alpha) - \tan p].\end{aligned}$$

5. The angle, ϕ , between two lines :


This is equal to the distance between their poles ; therefore,

$$\cos \phi = \sin p \sin p' + \cos p \cos p' \cos (\alpha' - \alpha).$$

The two lines $a \tan x + b \tan y + 1 = 0$,

$$a' \tan x + b' \tan y + 1 = 0$$

are perpendicular if $aa' + bb' + 1 = 0$.




6. Differential formulæ :

The formula
$$\frac{\sin A}{\sin a} = \frac{\sin C}{\sin c}$$

becomes, when A diminishes indefinitely,

$$(I.) \quad a = \sin c \cdot A.$$


Corollary. *In a circle of radius r , the ratio of any arc to the angle subtended at the centre is $\sin r$.*



From the right triangle ABC , if b remain fixed, we get, by differentiating the equation

$$\sin A = \frac{\cos B}{\cos b},$$

$$(II.) \quad dB = -\cos c \, dA.$$



Thus, we have for differential formulæ in polar coördinates

$$ds^2 = d\rho^2 + \sin^2 \rho \, d\theta^2,$$

$$\tan \phi = \sin \rho \frac{d\theta}{d\rho}; *$$

* If ϕ is constant, as in the logarithmic spiral of Euclidean Geometry, we can integrate this equation; namely,

$$\tan \phi \frac{d\rho}{\sin \rho} = d\theta.$$

$$\therefore \tan \phi \log \tan \frac{\rho}{2} = \theta + c,$$

or

$$\tan \frac{\rho}{2} = e^{\frac{\theta + c}{\tan \phi}}.$$

Writing c' for $\frac{c}{\tan \phi}$, this is

$$\tan \frac{\rho}{2} = c' e^{\frac{\theta}{\tan \phi}}.$$

On the sphere this is the curve called the loxodrome.

and for two arcs cutting at right angles

$$\frac{d\rho}{d\theta} \frac{d'\rho}{d'\theta} = -\sin^2\rho.$$

The formula for area is *

$$\iint \sin \rho \, d\rho \, d\theta.$$

We integrate first with respect to ρ , and if the area contains the origin and each radius vector meets the curve once, and only once, our expression becomes

$$2\pi - \int_0^{2\pi} \cos \rho \, d\theta.$$

The entire rotation in going around the curve is found as on page 79, and is

$$\int_0^{2\pi} \cos \rho \, d\theta.$$

Thus the area is equal to the amount by which this rotation is less than four right angles.

For example, the area of a circle of radius ρ is $2\pi(1 - \cos \rho)$, and the amount of turning in going around it is $2\pi \cos \rho$. The area of the entire plane is 2π .

7. A modified system of coördinates :

Writing u for $\tan x$, v for $\tan y$, r for $\tan \rho$, etc., we have

$$u^2 + v^2 = r^2. \dagger$$

The equation of a line then becomes

$$au + bv + 1 = 0,$$

and the equation of a circle


$$(1 + u^2 + v^2)c^2 = (1 + au + bv)^2.$$

* The unit of area being properly chosen.

† The footnote on page 80 applies here also.

III. ELLIPTIC SOLID ANALYTIC GEOMETRY

We will develop far enough to get the equation of the surface of double revolution.



1. Coördinates, lines, and planes :

Draw three planes through the point perpendicular to the axes. For coördinates x , y , z , we take the intercepts which these planes make on the axes.

The lines of intersection of these three planes are perpendicular to the coördinate planes (Chap. I, II, 16 and 17); in fact, all the face angles in the figure are right angles except those at P and the three angles $BA'C$, $CB'A$, and $AC'B$, which are obtuse angles.

Let ρ be the radius vector to the point P , and α , β , and γ the three angles which it makes with the three axes. Then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1,$$

$$\cos \alpha = \frac{\tan x}{\tan \rho}, \text{ etc.};$$

$$\tan^2 x + \tan^2 y + \tan^2 z = \tan^2 \rho.$$

For the angle between two lines intersecting at the origin

$$\cos \theta = \cos \alpha \cos \alpha' + \cos \beta \cos \beta' + \cos \gamma \cos \gamma'.$$

The angle subtended at the origin by the two points xyz and $x'y'z'$ is given by the equation

$$\cos \theta = \frac{\tan x \tan x' + \tan y \tan y' + \tan z \tan z'}{\tan \rho \tan \rho'}.$$

For the distance between two points

$$\cos \delta = \cos \rho \cos \rho' + \sin \rho \sin \rho' \cos \theta.$$

This gives us the equation of a sphere, and for $\delta = \frac{\pi}{2}$ the equation of a plane. The latter in rectangular coördinates is

$$\tan x \tan x' + \tan y \tan y' + \tan z \tan z' + 1 = 0.$$

Let p be the length of the perpendicular from the origin upon the plane, and α, β, γ the angles which this perpendicular makes with the axes. Then we have for its pole

$$\rho' = p + \frac{\pi}{2},$$

$$\tan x' = \tan \rho' \cos \alpha = -\frac{\cos \alpha}{\tan p}, \text{ etc.};$$

hence, the equation of the plane may be written

$$\cos \alpha \tan x + \cos \beta \tan y + \cos \gamma \tan z = \tan p.$$

2. The surface of double revolution:

Take one of its axes for the axis of z , suppose k the distance of the surface from this axis, and let θ denote the angle which the plane through the point P and the axis of z makes with the plane of xz . We may call z and θ latitude and longitude.

Produce OA and CB . They will meet at a distance, $\frac{\pi}{2}$, from the axis of z in a point, O' , on the other axis of the surface, and there form an angle that is equal in measure to z .

That is, in terms of z , z' , θ , and θ' , we have

$$\cos \delta = \cos^2 k \cos (z' - z) + \sin^2 k \cos (\theta' - \theta).$$

From this we can get an expression for ds , the differential element of length on the surface :

$$\cos ds = \cos^2 k \cos dz + \sin^2 k \cos d\theta,$$

or, since $\cos ds = 1 - \frac{ds^2}{2}$, etc.,

$$ds^2 = \cos^2 k dz^2 + \sin^2 k d\theta^2.$$

z and θ are proportional to the distances measured along the two systems of circles. These circles cut at right angles, and may be used to give us a system of rectangular coördinates on the surface. The actual lengths along these two systems of circles are $\theta \sin k$ and $z \cos k$ (see Cor. p. 83). If, therefore, we write

$$\alpha = \theta \sin k, \quad \beta = z \cos k,$$

we shall have a rectangular system on the surface where the coördinates are the distances measured along these two systems of circles which cut at right angles.

The formula now becomes

$$ds^2 = d\alpha^2 + d\beta^2.$$

An equation of the first degree in α and β represents a curve which enjoys on this surface all the properties of the straight line in the plane of the Euclidean Geometry. Through any two points one, and only one, such line can be drawn, because two sets of coördinates are just sufficient to determine the coefficients of an equation of the first degree. The shortest distance between two points on the surface is measured on such a line. For, the distance between two points on a path represented by an equation in α and β is the same as the distance between the corresponding points and on the corresponding path in a Euclidean plane in which we take α and β for rectangular coördinates. It must, therefore, be the shortest

when the path is represented by an equation of the first degree in α and β . Such a line on a surface is called a *geodesic line*, or, so far as the surface is concerned, a straight line. The distance between any two points measured on one of these lines is expressed by the formula

$$d = \sqrt{(\alpha - \alpha')^2 + (\beta - \beta')^2}.$$

Triangles formed of these lines have all the properties of plane triangles in the Euclidean Geometry: the sum of the angles is π , etc. In fact this surface has the same relation to elliptic space that the boundary-surface has to hyperbolic space.

The normal form of the equation of a line is

$$\alpha \cos \omega + \beta \sin \omega = p.$$

The rectilinear generators of the surface make a constant angle, $\pm k$, with all the circles drawn around the axis which is polar to the axis of z . These generators are then "straight lines" on the surface, and their equation takes the form

$$\alpha \cos k \pm \beta \sin k = p.$$

HISTORICAL NOTE

THE history of Non-Euclidean Geometry has been so well and so often written that we will give only a brief outline.

There is one axiom of Euclid that is somewhat complicated in its expression and does not seem to be, like the rest, a simple elementary fact. It is this:*

If two lines are cut by a third, and the sum of the interior angles on the same side of the cutting line is less than two right angles, the lines will meet on that side when sufficiently produced.

Attempts were made by many mathematicians, notably by Legendre, to give a proof of this proposition; that is, to show that it is a necessary consequence of the simpler axioms preceding it. Legendre proved that the sum of the angles of a triangle can never exceed two right angles, and that if there is a single triangle in which this sum is equal to two right angles, the same is true of all triangles. This was, of course, on the supposition that a line is of infinite length. He could not, however, prove that there exists a triangle the sum of whose angles is two right angles.†

At last some mathematicians began to believe that this statement was not capable of proof, that an equally consistent

* See article on the axioms of Euclid by Paul Tannery, *Bulletin des Sciences Mathématiques*, 1884.

† See, for example, the twelfth edition of his *Éléments de Géométrie*, Livre I, Proposition XIX, and Note II. See also a statement by Klein in an article on the Non-Euclidean Geometry in the second volume of the first series of the *Bulletin des Sciences Mathématiques*.

Geometry could be built up if we suppose it not always true, and, finally, that all of the postulates of Euclid were only hypotheses which our experience had led us to accept as true, but which could be replaced by contrary statements in the development of a logical Geometry.

The beginnings of this theory have sometimes been ascribed to Gauss, but it is known now that a paper was written by Lambert,* in 1766, in which he maintains that the parallel axiom needs proof, and gives some of the characteristics of Geometries in which this axiom does not hold. Even as long ago as 1733 a book was published by an Italian, Saccheri, in which he gives a complete system of Non-Euclidean Geometry, and then saves himself and his book by asserting dogmatically that these other hypotheses are false. It is his method of treatment that has been taken as the basis of the first chapter of this book.†

Gauss was seeking to prove the axiom of parallels for many years, and he may have discovered some of the theorems which are consequences of the denial of this axiom, but he never published anything on the subject.

Lobachevsky, in Russia, and Johann Bolyai, in Hungary, first asserted and proved that the axiom of parallels is not necessarily true. They were entirely independent of each other in their work, and each is entitled to the full credit of this discovery. Their results were published about 1830.

It was a long time before these discoveries attracted much notice. Meanwhile, other lines of investigation were carried on which were afterwards to throw much light on our subject, not, indeed, as explanations, but by their striking analogies.

Thus, within a year or two of each other, in the same journal (Crelle) appeared an article by Lobachevsky giving

* See *American Mathematical Monthly*, July-August, 1895.

† The translation of Saccheri by Halsted has been appearing in the *American Mathematical Monthly*.

the results of his investigations, and a memoir by Minding on surfaces on which he found that the formulæ of Spherical Trigonometry hold if we put ia for a , etc. Yet these two papers had been published thirty years before their connection was noticed (by Beltrami).

Again, Cayley, in 1859, in the *Philosophical Transactions*, published his Sixth Memoir on Quantics, in which he developed a projective theory of measurement and showed how metrical properties can be treated as projective by considering the anharmonic relations of any figures with a certain special figure that he called the absolute. In 1872 Klein took up this theory and showed that it gave a perfect image of the Non-Euclidean Geometry.

It has also been shown that we can get our Non-Euclidean Geometries if we think of a unit of measure varying according to a certain law as it moves about in a plane or in space.

The older workers in these fields discovered only the Geometry in which the hypothesis of the acute angle is assumed. It did not occur to them to investigate the assumption that a line is of finite length. The Elliptic Geometry was left to be discovered by Riemann, who, in 1854, took up a study of the foundations of Geometry. He studied it from a very different point of view, an abstract algebraic point of view, considering not our space and geometrical figures, except by way of illustration, but a system of variables. He investigated the question, What is the nature of a function of these variables which can be called element of length or distance? and found that in the simplest cases it must be the square root of a quadratic function of the differentials of the variables whose coefficients may themselves be functions of the variables. By taking different forms of the quadratic expressions we get an infinite number of these different kinds of Geometry, but in most of them we lose the axiom that bodies may be moved about without changing their size or shape.

Two more names should be included in this sketch, — Helmholtz and Clifford. These did much to make the subject popular by articles in scientific journals. To Clifford we owe the theory of parallels in elliptic space, as explained on page 68. He showed that we can have in this Geometry a finite surface on which the Euclidean Geometry holds true.*

The chief lesson of Non-Euclidean Geometry is that the axioms of Geometry are only deductions from our experience, like the theories of physical science. For the mathematician, they are hypotheses whose truth or falsity does not concern him, but only the philosopher. He may take them in any form he pleases and on them build his Geometry, and the Geometries so obtained have their applications in other branches of mathematics.

The "axiom," so far as this word is applied to these geometrical propositions, is not "self-evident," and is not necessarily true. If a certain statement can be proved, — that is, if it is a necessary consequence of axioms already adopted, — then it should not be called an axiom. When two or more mutually contradictory statements are equally consistent with all the axioms that have already been accepted, then we are at liberty to take either of them, and the statement which we choose

* Some of the more interesting accounts of Non-Euclidean Geometry are: *Encyclopedia Britannica*, article "Measurement," by Sir Robert Ball. *Revue Générale des Sciences*, 1891, "Les Géométries Non-Euclidean," by Poincaré. *Bulletin of the American Mathematical Society*, May and June, 1900, "Lobachevsky's Geometry," by Frederick S. Woods. *Mathematische Annalen*, Bd. xlix, p. 149, 1897, and *Bulletin des Sciences Mathématiques*, 1897, "Letters of Gauss and Bolyai"; particularly interesting is one letter in which Gauss gives a formula for the area of a triangle on the hypothesis that we can draw three mutually parallel lines enclosing a finite area always the same. The last two articles refer to the publications of Professors Engel and Stäckel, which give in German a full history of the theory of parallels and the writings and lives of Lobachevsky and Bolyai. See also the translations by Prof. George Bruce Halsted of Lobachevsky and Bolyai and of an address by Professor Vasiliev.

becomes for our Geometry an axiom. Our Geometry is a study of the consequences of this axiom.

The assumptions which distinguish the three kinds of Geometry that we have been studying may be expressed in different forms. We may say that one or two or no parallels can be drawn through a point; or, that the sum of the angles of a triangle is equal to, less than, or greater than two right angles; or, that a straight line has two real points, one real point, or no real point at infinity; or, that in a plane we can have similar figures or we cannot have similar figures, and a straight line is of finite or infinite length, etc. But any of these forms determines the nature of the Geometry, and the others are deducible from it.

CATALOG OF DOVER BOOKS

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

THE COMMON SENSE OF THE EXACT SCIENCES, W. K. Clifford. Introduction by James Newman, edited by Karl Pearson. For 70 years this has been a guide to classical scientific and mathematical thought. Explains with unusual clarity basic concepts, such as extension of meaning of symbols, characteristics of surface boundaries, properties of plane figures, vectors, Cartesian method of determining position, etc. Long preface by Bertrand Russell. Bibliography of Clifford. Corrected, 130 diagrams redrawn. 249pp. 5½ x 8.

T61 Paperbound \$1.60

SCIENCE THEORY AND MAN, Erwin Schrödinger. This is a complete and unabridged reissue of *SCIENCE AND THE HUMAN TEMPERAMENT* plus an additional essay: "What is an Elementary Particle?" Nobel Laureate Schrödinger discusses such topics as nature of scientific method, the nature of science, chance and determinism, science and society, conceptual models for physical entities, elementary particles and wave mechanics. Presentation is popular and may be followed by most people with little or no scientific training. "Fine practical preparation for a time when laws of nature, human institutions . . . are undergoing a critical examination without parallel," Waldemar Kaempffert, N. Y. TIMES. 192pp. 5½ x 8.

T428 Paperbound \$1.35

PIONEERS OF SCIENCE, O. Lodge. Eminent scientist-expositor's authoritative, yet elementary survey of great scientific theories. Concentrating on individuals—Copernicus, Brahe, Kepler, Galileo, Descartes, Newton, Laplace, Herschel, Lord Kelvin, and other scientists—the author presents their discoveries in historical order adding biographical material on each man and full, specific explanations of their achievements. The clear and complete treatment of the post-Newtonian astronomers is a feature seldom found in other books on the subject. Index. 120 illustrations. xv + 404pp. 5½ x 8.

T716 Paperbound \$1.50

THE EVOLUTION OF SCIENTIFIC THOUGHT FROM NEWTON TO EINSTEIN, A. d'Abro. Einstein's special and general theories of relativity, with their historical implications, are analyzed in non-technical terms. Excellent accounts of the contributions of Newton, Riemann, Weyl, Planck, Eddington, Maxwell, Lorentz and others are treated in terms of space and time, equations of electromagnetics, finiteness of the universe, methodology of science. 21 diagrams. 482pp. 5½ x 8.

T2 Paperbound \$2.00

THE RISE OF THE NEW PHYSICS, A. d'Abro. A half-million word exposition, formerly titled *THE DECLINE OF MECHANISM*, for readers not versed in higher mathematics. The only thorough explanation, in everyday language, of the central core of modern mathematical physical theory, treating both classical and modern theoretical physics, and presenting in terms almost anyone can understand the equivalent of 5 years of study of mathematical physics. Scientifically impeccable coverage of mathematical-physical thought from the Newtonian system up through the electronic theories of Dirac and Heisenberg and Fermi's statistics. Combines both history and exposition; provides a broad yet unified and detailed view, with constant comparison of classical and modern views on phenomena and theories. "A must for anyone doing serious study in the physical sciences," JOURNAL OF THE FRANKLIN INSTITUTE. "Extraordinary faculty . . . to explain ideas and theories of theoretical physics in the language of daily life," ISIS. First part of set covers philosophy of science, drawing upon the practice of Newton, Maxwell, Poincaré, Einstein, others, discussing modes of thought, experiment, interpretations of causality, etc. In the second part, 100 pages explain grammar and vocabulary of mathematics, with discussions of functions, groups, series, Fourier series, etc. The remainder is devoted to concrete, detailed coverage of both classical and quantum physics, explaining such topics as analytic mechanics, Hamilton's principle, wave theory of light, electromagnetic waves, groups of transformations, thermodynamics, phase rule, Brownian movement, kinetics, special relativity, Planck's original quantum theory, Bohr's atom, Zeeman effect, Broglie's wave mechanics, Heisenberg's uncertainty, Eigen-values, matrices, scores of other important topics. Discoveries and theories are covered for such men as Alembert, Born, Cantor, Debye, Euler, Foucault, Galois, Gauss, Hadamard, Kelvin, Kepler, Laplace, Maxwell, Pauli, Rayleigh, Volterra, Weyl, Young, more than 180 others. Indexed. 97 illustrations. ix + 982pp. 5½ x 8.

T3 Volume 1, Paperbound \$2.00

T4 Volume 2, Paperbound \$2.00

CONCERNING THE NATURE OF THINGS, Sir William Bragg. Christmas lectures delivered at the Royal Society by Nobel laureate. Why a spinning ball travels in a curved track; how uranium is transmuted to lead, etc. Partial contents: atoms, gases, liquids, crystals, metals, etc. No scientific background needed; wonderful for intelligent child. 32pp. of photos, 57 figures. xii + 232pp. 5½ x 8.

T31 Paperbound \$1.35

THE UNIVERSE OF LIGHT, Sir William Bragg. No scientific training needed to read Nobel Prize winner's expansion of his Royal Institute Christmas Lectures. Insight into nature of light, methods and philosophy of science. Explains lenses, reflection, color, resonance, polarization, x-rays, the spectrum, Newton's work with prisms, Huygens' with polarization, Crookes' with cathode ray, etc. Leads into clear statement of 2 major historical theories of light, corpuscle and wave. Dozens of experiments you can do. 199 illus., including 2 full-page color plates. 293pp. 5½ x 8.

S538 Paperbound \$1.85

BRIDGES AND THEIR BUILDERS, David Steinman and Sara Ruth Watson. Engineers, historians, everyone who has ever been fascinated by great spans will find this book an endless source of information and interest. Dr. Steinman, recipient of the Louis Levy medal, was one of the great bridge architects and engineers of all time, and his analysis of the great bridges of history is both authoritative and easily followed. Greek and Roman bridges, medieval bridges, Oriental bridges, modern works such as the Brooklyn Bridge and the Golden Gate Bridge, and many others are described in terms of history, constructional principles, artistry, and function. All in all this book is the most comprehensive and accurate semipopular history of bridges in print in English. New, greatly revised, enlarged edition. 23 photographs, 26 line drawings. Index. xvii + 401pp. 5½ x 8. T431 Paperbound \$2.00

FADS AND FALLACIES IN THE NAME OF SCIENCE, Martin Gardner. Examines various cults, quack systems, frauds, delusions which at various times have masqueraded as science. Accounts of hollow-earth fanatics like Symmes; Velikovsky and wandering planets; Hoerbiger; Bellamy and the theory of multiple moons; Charles Fort; dowsing, pseudoscientific methods for finding water, ores, oil. Sections on naturopathy, iridagnosis, zone therapy, food fads, etc. Analytical accounts of Wilhelm Reich and orgone sex energy; L. Ron Hubbard and Dianetics; A. Korzybski and General Semantics; many others. Brought up to date to include Bridey Murphy, others. Not just a collection of anecdotes, but a fair, reasoned appraisal of eccentric theory. Formerly titled *IN THE NAME OF SCIENCE*. Preface. Index. x + 384pp. 5½ x 8. T394 Paperbound \$1.50

See also: **A PHILOSOPHICAL ESSAY ON PROBABILITIES**, P. de Laplace; **ON MATHEMATICS AND MATHEMATICIANS**, R. E. Moritz; **AN ELEMENTARY SURVEY OF CELESTIAL MECHANICS**, Y. Ryabov; **THE SKY AND ITS MYSTERIES**, E. A. Beetz; **THE REALM OF THE NEBULAE**, E. Hubble; **OUT OF THE SKY**, H. H. Nininger; **SATELLITES AND SCIENTIFIC RESEARCH**, D. King-Hele; **HEREDITY AND YOUR LIFE**, A. M. Winchester; **INSECTS AND INSECT LIFE**, S. W. Frost; **PRINCIPLES OF STRATIGRAPHY**, A. W. Grabau; **TEACH YOURSELF SERIES**.

HISTORY OF SCIENCE AND MATHEMATICS

DIALOGUES CONCERNING TWO NEW SCIENCES, Galileo Galilei. This classic of experimental science, mechanics, engineering, is as enjoyable as it is important. A great historical document giving insights into one of the world's most original thinkers, it is based on 30 years' experimentation. It offers a lively exposition of dynamics, elasticity, sound, ballistics, strength of materials, the scientific method. "Superior to everything else of mine," Galileo. Trans. by H. Crew, A. Salvo. 126 diagrams. Index. xxi + 288pp. 5½ x 8. S99 Paperbound \$1.65

A DIDEROT PICTORIAL ENCYCLOPEDIA OF TRADES AND INDUSTRY, Manufacturing and the Technical Arts in Plates Selected from "L'Encyclopedie ou Dictionnaire Raisonné des Sciences, des Arts, et des Métiers" of Denis Diderot. Edited with text by C. Gillispie. This first modern selection of plates from the high point of 18th century French engraving is a storehouse of valuable technological information to the historian of arts and science. Over 2000 illustrations on 485 full page plates, most of them original size, show the trades and industries of a fascinating era in such great detail that the processes and shops might very well be reconstructed from them. The plates teem with life, with men, women, and children performing all of the thousands of operations necessary to the trades before and during the early stages of the industrial revolution. Plates are in sequence, and show general operations, closeups of difficult operations, and details of complex machinery. Such important and interesting trades and industries are illustrated as sowing, harvesting, bee-keeping, cheesemaking, operating windmills, milling flour, charcoal burning, tobacco processing, indigo, fishing, arts of war, salt extraction, mining, smelting, casting iron, steel, extracting mercury, zinc, sulphur, copper, etc., slating, tinning, silverplating, gilding, making gunpowder, cannons, bells, shoeing horses, tanning, papermaking, printing, dyeing, and more than 40 other categories. Professor Gillispie, of Princeton, supplies a full commentary on all the plates, identifying operations, tools, processes, etc. This material, presented in a lively and lucid fashion, is of great interest to the reader interested in history of science and technology. Heavy library cloth. 920pp. 9 x 12. T421 Two volume set \$18.50

DE MAGNETE, William Gilbert. This classic work on magnetism founded a new science. Gilbert was the first to use the word "electricity", to recognize mass as distinct from weight, to discover the effect of heat on magnetic bodies; invent an electroscope, differentiate between static electricity and magnetism, conceive of the earth as a magnet. Written by the first great experimental scientist, this lively work is valuable not only as an historical landmark, but as the delightfully easy to follow record of a perpetually searching, ingenious mind. Translated by P. F. Mottelay. 25 page biographical memoir. 90 figures. lix + 368pp. 5½ x 8. S470 Paperbound \$2.00

CHARLES BABBAGE AND HIS CALCULATING ENGINES, edited by P. Morrison and E. Morrison. Babbage, leading 19th century pioneer in mathematical machines and herald of modern operational research, was the true father of Harvard's relay computer Mark I. His Difference Engine and Analytical Engine were the first machines in the field. This volume contains a valuable introduction on his life and work; major excerpts from his autobiography, revealing his eccentric and unusual personality; and extensive selections from "Babbage's Calculating Engines," a compilation of hard-to-find journal articles by Babbage, the Countess of Lovelace, L. F. Menabrea, and Dionysius Lardner. 8 illustrations, Appendix of miscellaneous papers. Index. Bibliography. xxxviii + 400pp. 5½ x 8. T12 Paperbound \$2.00

A HISTORY OF ASTRONOMY FROM THALES TO KEPLER, J. L. E. Dreyer. (Formerly A HISTORY OF PLANETARY SYSTEMS FROM THALES TO KEPLER.) This is the only work in English to give the complete history of man's cosmological views from prehistoric times to Kepler and Newton. Partial contents: Near Eastern astronomical systems, Early Greeks, Homocentric Spheres of Eudoxus, Epicycles, Ptolemaic system, medieval cosmology, Copernicus, Kepler, etc. Revised, foreword by W. H. Stahl. New bibliography. xvii + 430pp. 5½ x 8. S79 Paperbound \$1.98

A SHORT HISTORY OF ANATOMY AND PHYSIOLOGY FROM THE GREEKS TO HARVEY, Charles Singer. Corrected edition of THE EVOLUTION OF ANATOMY, classic work tracing evolution of anatomy and physiology from prescientific times through Greek & Roman periods, Dark Ages, Renaissance, to age of Harvey and beginning of modern concepts. Centered on individuals, movements, periods that definitely advanced anatomical knowledge: Plato, Diocles, Aristotle, Theophrastus, Herophilus, Erasistratus, the Alexandrians, Galen, Mondino, da Vinci, Linacre, Sylvius, others. Special section on Vesalius; Vesalian atlas of nudes, skeletons, muscle tabulae. Index of names, 20 plates. 270 extremely interesting illustrations of ancient, medieval, Renaissance, Oriental origin. xii + 209pp. 5½ x 8. T389 Paperbound \$1.75

FROM MAGIC TO SCIENCE, Charles Singer. A great historian examines aspects of medical science from the Roman Empire through the Renaissance. Includes perhaps the best discussion of early herbals, and a penetrating physiological interpretation of "The Visions of Hildegard of Bingen." Also examined are Arabian and Galenic influences; the Sphere of Pythagoras; Paracelsus; the reawakening of science under Leonardo da Vinci, Vesalius; the Lorica of Gildas the Briton; etc. Frequent quotations with translations. New Introduction by the author. New unabridged, corrected edition. 158 unusual illustrations from classical and medieval sources. Index. xxvii + 365pp. 5½ x 8. T390 Paperbound \$2.00

HISTORY OF MATHEMATICS, D. E. Smith. Most comprehensive non-technical history of math in English. Discusses lives and works of over a thousand major and minor figures, with footnotes supplying technical information outside the book's scheme, and indicating disputed matters. Vol I: A chronological examination, from primitive concepts through Egypt, Babylonia, Greece, the Orient, Rome, the Middle Ages, the Renaissance, and up to 1900. Vol 2: The development of ideas in specific fields and problems, up through elementary calculus. Two volumes, total of 510 illustrations, 1355pp. 5½ x 8. Set boxed in attractive container. T429, 430 Paperbound, the set \$5.00

A SHORT ACCOUNT OF THE HISTORY OF MATHEMATICS, W. W. R. Ball. Most readable non-technical history of mathematics treats lives, discoveries of every important figure from Egyptian, Phoenician mathematicians to late 19th century. Discusses schools of Ionia, Pythagoras, Athens, Cyzicus, Alexandria, Byzantium, systems of numeration; primitive arithmetic; Middle Ages, Renaissance, including Arabs, Bacon, Regiomontanus, Tartaglia, Cardan, Stevinus, Galileo, Kepler; modern mathematics of Descartes, Pascal, Wallis, Huygens, Newton, Leibnitz, d'Alembert, Euler, Lambert, Laplace, Legendre, Gauss, Hermite, Weierstrass, scores more. Index. 25 figures. 546pp. 5½ x 8. S630 Paperbound \$2.00

A SOURCE BOOK IN MATHEMATICS, D. E. Smith. Great discoveries in math, from Renaissance to end of 19th century, in English translation. Read announcements by Dedekind, Gauss, Delamain, Pascal, Fermat, Newton, Abel, Lobachevsky, Bolyai, Riemann, De Moivre, Legendre, Laplace, others of discoveries about imaginary numbers, number congruence, slide rule, equations, symbolism, cubic algebraic equations, non-Euclidean forms of geometry, calculus, function theory, quaternions, etc. Succinct selections from 125 different treatises, articles, most unavailable elsewhere in English. Each article preceded by biographical, historical introduction. Vol. I: Fields of Number, Algebra. Index. 32 illus. 338pp. 5½ x 8. Vol. II: Fields of Geometry, Probability, Calculus, Functions, Quaternions. 83 illus. 432pp. 5½ x 8. Vol. 1: \$552 Paperbound \$1.85 Vol. 2: \$553 Paperbound \$1.85 2 vol. set, boxed \$3.50

A HISTORY OF THE CALCULUS, AND ITS CONCEPTUAL DEVELOPMENT, Carl B. Boyer. Provides laymen and mathematicians a detailed history of the development of the calculus, from early beginning in antiquity to final elaboration as mathematical abstractions. Gives a sense of mathematics not as a technique, but as a habit of mind, in the progression of ideas of Zeno, Plato, Pythagoras, Eudoxus, Arabic and Scholastic mathematicians, Newton, Leibnitz, Taylor, Descartes, Euler, Lagrange, Cantor, Weierstrass, and others. This first comprehensive critical history of the calculus was originally titled "The Concepts of the Calculus." Foreword by R. Courant. Preface. 22 figures. 25-page bibliography. Index. v + 364pp. 5½ x 8. S509 Paperbound \$2.00

A CONCISE HISTORY OF MATHEMATICS, D. Struik. Lucid study of development of mathematical ideas, techniques from Ancient Near East, Greece, Islamic science, Middle Ages, Renaissance, modern times. Important mathematicians are described in detail. Treatment is not anecdotal, but analytical development of ideas. "Rich in content, thoughtful in interpretation," U.S. QUARTERLY BOOKLIST. Non-technical; no mathematical training needed. Index. 60 illustrations, including Egyptian papyri, Greek mss., portraits of 31 eminent mathematicians. Bibliography. 2nd edition. xix + 299pp. 5½ x 8. T255 Paperbound \$1.75

See also: **NON-EUCLIDEAN GEOMETRY, R. Bonola; THEORY OF DETERMINANTS IN HISTORICAL ORDER OF DEVELOPMENT, T. Muir; HISTORY OF THE THEORY OF ELASTICITY AND STRENGTH OF MATERIALS, I. Todhunter and K. Pearson; A SHORT HISTORY OF ASTRONOMY, A. Berry; CLASSICS OF SCIENCE.**

PHILOSOPHY OF SCIENCE AND MATHEMATICS

FOUNDATIONS OF SCIENCE: THE PHILOSOPHY OF THEORY AND EXPERIMENT, N. R. Campbell. A critique of the most fundamental concepts of science in general and physics in particular. Examines why certain propositions are accepted without question, demarcates science from philosophy, clarifies the understanding of the tools of science. Part One analyzes the presuppositions of scientific thought: existence of the material world, nature of scientific laws, multiplication of probabilities, etc. Part Two covers the nature of experiment and the application of mathematics: conditions for measurement, relations between numerical laws and theories, laws of error, etc. An appendix covers problems arising from relativity, force, motion, space, and time. A classic in its field. Index. xiii + 565pp. 5½ x 8. S372 Paperbound \$2.95

WHAT IS SCIENCE?, Norman Campbell. This excellent introduction explains scientific method, role of mathematics, types of scientific laws. Contents: 2 aspects of science, science & nature, laws of science, discovery of laws, explanation of laws, measurement & numerical laws, applications of science. 192pp. 5½ x 8. S43 Paperbound \$1.25

THE VALUE OF SCIENCE, Henri Poincaré. Many of the most mature ideas of the "last scientific universalist" covered with charm and vigor for both the beginning student and the advanced worker. Discusses the nature of scientific truth, whether order is innate in the universe or imposed upon it by man, logical thought versus intuition (relating to math, through the works of Weierstrass, Lie, Klein, Riemann), time and space (relativity, psychological time, simultaneity), Hertz's concept of force, interrelationship of mathematical physics to pure math, values within disciplines of Maxwell, Carnot, Mayer, Newton, Lorentz, etc. Index. iii + 147pp. 5½ x 8. S469 Paperbound \$1.35

SCIENCE AND METHOD, Henri Poincaré. Procedure of scientific discovery, methodology, experiment, idea-germination—the intellectual processes by which discoveries come into being. Most significant and most interesting aspects of development, application of ideas. Chapters cover selection of facts, chance, mathematical reasoning, mathematics, and logic; Whitehead, Russell, Cantor; the new mechanics, etc. 288pp. 5½ x 8. S222 Paperbound \$1.35

SCIENCE AND HYPOTHESIS, Henri Poincaré. Creative psychology in science. How such concepts as number, magnitude, space, force, classical mechanics were developed, and how the modern scientist uses them in his thought. Hypothesis in physics, theories of modern physics. Introduction by Sir James Larmor. "Few mathematicians have had the breadth of vision of Poincaré, and none is his superior in the gift of clear exposition," E. T. Bell. Index. 272pp. 5½ x 8. S221 Paperbound \$1.35

PHILOSOPHY AND THE PHYSICISTS, L. S. Stebbing. The philosophical aspects of modern science examined in terms of a lively critical attack on the ideas of Jeans and Eddington. Discusses the task of science, causality, determinism, probability, consciousness, the relation of the world of physics to that of everyday experience. Probes the philosophical significance of the Planck-Bohr concept of discontinuous energy levels, the inferences to be drawn from Heisenberg's Uncertainty Principle, the implications of "becoming" involved in the 2nd law of thermodynamics, and other problems posed by the discarding of Laplacean determinism. 285pp. 5½ x 8. T480 Paperbound \$1.65

EXPERIMENT AND THEORY IN PHYSICS, Max Born. A Nobel laureate examines the nature and value of the counterclaims of experiment and theory in physics. Synthetic versus analytical scientific advances are analyzed in the work of Einstein, Bohr, Heisenberg, Planck, Eddington, Milne, and others by a fellow participant. 44nn. 5½ x 8. S222 Paperbound \$1.35

THE NATURE OF PHYSICAL THEORY, P. W. Bridgman. Here is how modern physics looks to a highly unorthodox physicist—a Nobel laureate. Pointing out many absurdities of science, and demonstrating the inadequacies of various physical theories, Dr. Bridgman weighs and analyzes the contributions of Einstein, Bohr, Newton, Heisenberg, and many others. This is a non-technical consideration of the correlation of science and reality. Index. xi + 138pp. 5½ x 8. S33 Paperbound \$1.25

THE PHILOSOPHY OF SPACE AND TIME, H. Reichenbach. An important landmark in the development of the empiricist conception of geometry, covering the problem of the foundations of geometry, the theory of time, the consequences of Einstein's relativity, including: relations between theory and observations; coordinate and metrical properties of space; the psychological problem of visual intuition of non-Euclidean structures; and many other important topics in modern science and philosophy. The majority of ideas require only a knowledge of intermediate math. Introduction by R. Carnap. 49 figures. Index. xviii + 296pp. 5½ x 8. S443 Paperbound \$2.00

MATTER & MOTION, James Clerk Maxwell. This excellent exposition begins with simple particles and proceeds gradually to physical systems beyond complete analysis: motion, force, properties of centre of mass of material system, work, energy, gravitation, etc. Written with all Maxwell's original insights and clarity. Notes by E. Larmor. 17 diagrams. 178pp. 5½ x 8. S188 Paperbound \$1.35

THE ANALYSIS OF MATTER, Bertrand Russell. How do our senses concord with the new physics? This volume covers such topics as logical analysis of physics, pre-relativity physics, causality, scientific inference, physics and perception, special and general relativity, Weyl's theory, tensors, invariants and their physical interpretation, periodicity and qualitative series. "The most thorough treatment of the subject that has yet been published," THE NATION. Introduction by L. E. Denonn. 422pp. 5½ x 8. T231 Paperbound \$1.95

SUBSTANCE AND FUNCTION, & EINSTEIN'S THEORY OF RELATIVITY, Ernst Cassirer. Two books bound as one. Cassirer establishes a philosophy of the exact sciences that takes into consideration newer developments in mathematics, and also shows historical connections. Partial contents: Aristotelian logic, Mill's analysis, Helmholtz & Kronecker, Russell & cardinal numbers, Euclidean vs. non-Euclidean geometry, Einstein's relativity. Bibliography. Index. xxi + 465pp. 5½ x 8. T50 Paperbound \$2.00

PRINCIPLES OF MECHANICS, Heinrich Hertz. This last work by the great 19th century physicist is not only a classic, but of great interest in the logic of science. Creating a new system of mechanics based upon space, time, and mass, it returns to axiomatic analysis, to understanding of the formal or structural aspects of science, taking into account logic, observation, and a priori elements. Of great historical importance to Poincaré, Carnap, Einstein, Milne. A 20-page introduction by R. S. Cohen, Wesleyan University, analyzes the implications of Hertz's thought and the logic of science. Bibliography. 13-page introduction by Helmholtz. xiii + 274pp. 5½ x 8. S316 Clothbound \$3.50
S317 Paperbound \$1.85

THE PHILOSOPHICAL WRITINGS OF PEIRCE, edited by Justus Buchler. (Formerly published as THE PHILOSOPHY OF PEIRCE.) This is a carefully balanced exposition of Peirce's complete system, written by Peirce himself. It covers such matters as scientific method, pure chance vs. law, symbolic logic, theory of signs, pragmatism, experiment, and other topics. Introduction by Justus Buchler, Columbia University. xvi + 368pp. 5½ x 8. T217 Paperbound \$1.95

ESSAYS IN EXPERIMENTAL LOGIC, John Dewey. This stimulating series of essays touches upon the relationship between inquiry and experience, dependence of knowledge upon thought, character of logic; judgments of practice, data and meanings, stimuli of thought, etc. Index. viii + 444pp. 5½ x 8. T73 Paperbound \$1.95

LANGUAGE, TRUTH AND LOGIC, A. Ayer. A clear introduction to the Vienna and Cambridge schools of Logical Positivism. It sets up specific tests by which you can evaluate validity of ideas, etc. Contents: Function of philosophy, elimination of metaphysics, nature of analysis; a priori, truth and probability, etc. 10th printing. "I should like to have written it myself," Bertrand Russell. Index. 160pp. 5½ x 8. T10 Paperbound \$1.25

THE PSYCHOLOGY OF INVENTION IN THE MATHEMATICAL FIELD, J. Hadamard. Where do ideas come from? What role does the unconscious play? Are ideas best developed by mathematical reasoning, word reasoning, visualization? What are the methods used by Einstein, Poincaré, Galton, Riemann? How can these techniques be applied by others? Hadamard, one of the world's leading mathematicians, discusses these and other questions. xiii + 145pp. 5½ x 8. T107 Paperbound \$1.25

FOUNDATIONS OF GEOMETRY, Bertrand Russell. Analyzing basic problems in the overlap area between mathematics and philosophy, Nobel laureate Russell examines the nature of geometrical knowledge, the nature of geometry, and the application of geometry to space. It covers the history of non-Euclidean geometry, philosophic interpretations of geometry—especially Kant—projective and metrical geometry. This is most interesting as the solution offered in 1897 by a great mind to a problem still current. New introduction by Prof. Morris Kline of N. Y. University. xii + 201pp. 5½ x 8. S232 Clothbound \$3.25
S233 Paperbound \$1.60

BIBLIOGRAPHIES

GUIDE TO THE LITERATURE OF MATHEMATICS AND PHYSICS, N. G. Parke III. Over 5000 entries included under approximately 120 major subject headings, of selected most important books, monographs, periodicals, articles in English, plus important works in German, French, Italian, Spanish, Russian (many recently available works). Covers every branch of physics, math, related engineering. Includes author, title, edition, publisher, place, date, number of volumes, number of pages. A 40-page introduction on the basic problems of research and study provides useful information on the organization and use of libraries, the psychology of learning, etc. This reference work will save you hours of time. 2nd revised edition. Indices of authors, subjects. 464pp. 5 3/8 x 8. \$447 Paperbound \$2.49

THE STUDY OF THE HISTORY OF MATHEMATICS & THE STUDY OF THE HISTORY OF SCIENCE, George Sarton. Scientific method & philosophy in 2 scholarly fields. Defines duty of historian of math., provides especially useful bibliography with best available biographies of modern mathematicians, editions of their collected works, correspondence. Observes combination of history & science, will aid scholar in understanding science today. Bibliography includes best known treatises on historical methods. 200-item critically evaluated bibliography. Index. 10 illustrations. 2 volumes bound as one. 113pp. + 75pp. 5 3/8 x 8. T240 Paperbound \$1.25

MATHEMATICAL PUZZLES

AMUSEMENTS IN MATHEMATICS, Henry Ernest Dudeney. The foremost British originator of mathematical puzzles is always intriguing, witty, and paradoxical in this classic, one of the largest collections of mathematical amusements. More than 430 puzzles, problems, and paradoxes. Mazes and games, problems on number manipulation, unicursal and other route problems, puzzles on measuring, weighing, packing, age, kinship, chessboards, joiners', crossing river, plane figure dissection, and many others. Solutions. More than 450 illustrations. vii + 258pp. 5 3/8 x 8. T473 Paperbound \$1.25

THE CANTERBURY PUZZLES, Henry Ernest Dudeney. Chaucer's pilgrims set one another problems in story form. Also Adventures of the Puzzle Club, the Strange Escape of the King's Jester, the Monks of Riddiwell, the Squire's Christmas Puzzle Party, and others. All puzzles are original, based on dissecting plane figures, arithmetic, algebra, elementary calculus, and other branches of mathematics, and purely logical ingenuity. "The limit of ingenuity and intricacy . . ." The Observer. Over 110 puzzles. Full solutions. 150 illustrations. vii + 225pp. 5 3/8 x 8. T474 Paperbound \$1.25

SYMBOLIC LOGIC and THE GAME OF LOGIC, Lewis Carroll. "Symbolic Logic" is not concerned with modern symbolic logic, but is instead a collection of over 380 problems posed with charm and imagination, using the syllogism, and a fascinating diagrammatic method of drawing conclusions. In "The Game of Logic," Carroll's whimsical imagination devises a logical game played with 2 diagrams and counters (included) to manipulate hundreds of tricky syllogisms. The final section, "Hit or Miss" is a lagniappe of 101 additional puzzles in the delightful Carroll manner. Until this reprint edition, both of these books were rarities costing up to \$15 each. Symbolic Logic: Index, xxxi + 199pp. The Game of Logic: 96pp. Two vols. bound as one. 5 3/8 x 8. T492 Paperbound \$1.50

PILLOW PROBLEMS and A TANGLED TALE, Lewis Carroll. One of the rarest of all Carroll's works, "Pillow Problems" contains 72 original math puzzles, all typically ingenious. Particularly fascinating are Carroll's answers which remain exactly as he thought them out, reflecting his actual mental processes. The problems in "A Tangled Tale" are in story form, originally appearing as a monthly magazine serial. Carroll not only gives the solutions, but uses answers sent in by readers to discuss wrong approaches and misleading paths, and grades them for insight. Both of these books were rarities until this edition, "Pillow Problems" costing up to \$25, and "A Tangled Tale" \$15. Pillow Problems: Preface and Introduction by Lewis Carroll. xx + 109pp. A Tangled Tale: 6 illustrations. 152pp. Two vols. bound as one. 5 3/8 x 8. T493 Paperbound \$1.50

DIVERSIONS AND DIGRESSIONS OF LEWIS CARROLL. A major new treasure for Carroll fans! Rare privately published puzzles, mathematical amusements and recreations, games. Includes the fragmentary Part III of "Curiosa Mathematica." Also contains humorous and satirical pieces: "The New Belfry," "The Vision of the Three T's," and much more. New 32-page supplement of rare photographs taken by Carroll. Formerly titled "The Lewis Carroll Picture Book." Edited by S. Collingwood. x + 37pp. 5 3/8 x 8. T494 Paperbound \$1.50

THE BOOK OF MODERN PUZZLES, G. L. Kaufman. More than 150 word puzzles, logic puzzles. No warmed-over fare but all new material based on same appeals that make crosswords and deduction puzzles popular, but with different principles, techniques. Two-minute teasers, involved word-labyrinths, design and pattern puzzles, puzzles calling for logic and observation, puzzles testing ability to apply general knowledge to peculiar situations, many others. Answers to all problems. 116 illustrations. 192pp. 5½ x 8. T143 Paperbound \$1.00

NEW WORD PUZZLES, Gerald L. Kaufman. Contains 100 brand new challenging puzzles based on words and their combinations, never published before in any form. Most are new types invented by the author—for beginners or experts. Chess word puzzles, addle letter anagrams, double word squares, double horizontals, alphagram puzzles, dual acrostigrams, linkogram lapwords—plus 8 other brand new types, all with solutions included. 196 figures. 100 brand new puzzles. vi + 122pp. 5½ x 8. T344 Paperbound \$1.00

MATHEMATICAL RECREATIONS

MATHEMATICS, MAGIC AND MYSTERY, Martin Gardner. Card tricks, feats of mental mathematics, stage mind-reading, other "magic" explained as applications of probability, sets, theory of numbers, topology, various branches of mathematics. Creative examination of laws and their applications with scores of new tricks and insights. 115 sections discuss tricks with cards, dice, coins; geometrical vanishing tricks, dozens of others. No sleight of hand needed; mathematics guarantees success. 115 illustrations. xii + 174pp. 5½ x 8. T335 Paperbound \$1.00

MATHEMATICAL EXCURSIONS, Helen A. Merrill. Fun, recreation, insights into elementary problem-solving. A mathematical expert guides you along by-paths not generally travelled in elementary math courses—how to divide by inspection, Russian peasant system of multiplication; memory systems for pi; building odd and even magic squares; dyadic systems; facts about 37; square roots by geometry; Tchebichev's machine; drawing five-sided figures; dozens more. Solutions to more difficult ones. 50 illustrations. 145pp. 5½ x 8. T350 Paperbound \$1.00

CRYPTOGRAPHY, L. D. Smith. Excellent elementary introduction to enciphering, deciphering secret writing. Explains transposition, substitution ciphers; codes; solutions. Geometrical patterns, route transcription, columnar transposition, other methods. Mixed cipher systems; single-alphabet, polyalphabetical substitution; mechanical devices; Vigenere system, etc. Enciphering Japanese; explanation of Baconian Biliteral cipher; frequency tables. More than 150 problems provide practical application. Bibliography. Index. 164pp. 5½ x 8. T247 Paperbound \$1.00

CRYPTANALYSIS, Helen F. Gaines. (Formerly ELEMENTARY CRYPTANALYSIS.) A standard elementary and intermediate text for serious students. It does not confine itself to old material, but contains much that is not generally known, except to experts. Concealment, Transposition, Substitution ciphers; Vigenere, Kasiski, Playfair, multifid, dozens of other techniques. Appendix with sequence charts, letter frequencies in English, 5 other languages, English word frequencies. Bibliography. 167 codes. New to this edition: solution to codes. vi + 230pp. 5½ x 8. T97 Paperbound \$1.95

MAGIC SQUARES AND CUBES, W. S. Andrews. Only book-length treatment in English, a thorough non-technical description and analysis. Here are nasik, overlapping, pandiagonal, serated squares; magic circles, cubes, spheres, rhombuses. Try your hand at 4-dimensional magical figures! Much unusual folklore and tradition included. High school algebra is sufficient. 754 diagrams and illustrations. viii + 419pp. 5½ x 8. T658 Paperbound \$1.85

PAPER FOLDING FOR BEGINNERS, W. D. Murray and F. J. Rigney. A delightful introduction to the varied and entertaining Japanese art of origami (paper folding), with a full crystal-clear text that anticipates every difficulty; over 275 clearly labeled diagrams of all important stages in creation. You get results at each stage, since complex figures are logically developed from simpler ones. 43 different pieces are explained: place mats, drinking cups, bonbon boxes, sailboats, frogs, roosters, etc. 6 photographic plates. 279 diagrams. 95pp. 5½ x 8½. T713 Paperbound \$1.00

CHESS, CHECKERS, GAMES, GO

A TREASURY OF CHESS LORE, edited by Fred Reinfeld. A delightful collection of anecdotes, short stories, aphorisms by and about the masters, poems, accounts of games and tournaments, photographs. Hundreds of humorous, pithy, satirical, wise, and historical episodes, comments, and word portraits. A fascinating "must" for chess players; revealing and perhaps seductive to those who wonder what their friends see in the game. 49 photographs (14 full page plates). 12 diagrams. xi + 306pp. 5½ x 8. T458 Paperbound \$1.75

HYPERMODERN CHESS as developed in the games of its greatest exponent, ARON NIMZOVICH, edited by Fred Reinfeld. An intensely original player and analyst, Nimzovich's extraordinary approaches startled and often angered the chess world. This volume, designed for the average player, shows in his victories over Alekhine, Lasker, Marshall, Rubinstein, Spielmann, and others, how his iconoclastic methods infused new life into the game. Use Nimzovich to invigorate your play and startle opponents. Introduction. Indices of players and openings. 180 diagrams. viii + 220pp. 5½ x 8. T448 Paperbound \$1.35

ONE HUNDRED SELECTED GAMES, Mikhail Botvinnik. Author's own choice of his best games before becoming World Champion in 1948, beginning with first big tournament, the USSR Championship, 1927. Shows his great powers of analysis as he annotates these games, giving strategy, technique against Alekhine, Capablanca, Euwe, Keres, Reshevsky, Smyslov, Vidmar, many others. Discusses his career, methods of play, system of training, 6 studies of endgame positions. 221 diagrams. 272pp. 5½ x 8. T620 Paperbound \$1.50

RUBINSTEIN'S CHESS MASTERPIECES, selected and annotated by H. Knoch. Thoroughgoing mastery of opening, middle game; faultless technique in endgame, particularly rook and pawn endings; ability to switch from careful positional play to daring combinations; all distinguish the play of Rubinstein. 100 best games, against Janowski, Nimzowitch, Tarrasch, Vidmar, Capablanca, other greats, carefully annotated, will improve your game rapidly. Biographical introduction, B. F. Winkelman, 103 diagrams. 192pp. 5½ x 8. T617 Paperbound \$1.25

TARRASCH'S BEST GAMES OF CHESS, selected & annotated by Fred Reinfeld. First definitive collection of games by Siegbert Tarrasch, winner of 7 international tournaments, and the leading theorist of classical chess. 183 games cover 50 years of play against Mason, Mieses, Paulsen, Teichmann, Pillsbury, Janowski, others. Reinfeld includes Tarrasch's own analyses of many of these games. A careful study and replaying of the games will give you a sound understanding of classical methods, and many hours of enjoyment. Introduction. Indexes. 183 diagrams. xxiv + 386pp. 5½ x 8. T644 Paperbound \$2.00

MARSHALL'S BEST GAMES OF CHESS, F. J. Marshall. Grandmaster, U. S. champion for 27 years, tells story of career; presents magnificent collection of 140 of best games, annotated by himself. Games against Capablanca, Alekhine, Emanuel Lasker, Janowski, Rubinstein, Pillsbury, etc. Special section analyzes openings such as King's Gambit, Ruy Lopez, Alekhine's Defence, Giuoco Piano, others. A study of Marshall's brilliant "swindles," slashing attacks, extraordinary sacrifices, will rapidly improve your game. Formerly "My Fifty Years of Chess." Introduction. 19 diagrams. 13 photos. 250pp. 5½ x 8. T604 Paperbound \$1.35

THE ENJOYMENT OF CHESS PROBLEMS, K. S. Howard. A classic treatise on this minor art by an internationally recognized authority that gives a basic knowledge of terms and themes for the everyday chess player as well as the problem fan: 7 chapters on the two-move; 7 more on 3- and 4-move problems; a chapter on selfmates; and much more. "The most important one-volume contribution originating solely in the U.S.A.," Alain White, 200 diagrams. Index. Solutions. viii + 212pp. 5½ x 8. T742 Paperbound \$1.25

HOW TO SOLVE CHESS PROBLEMS, K. S. Howard. Full of practical suggestions for the fan or the beginner—who need only know the moves of the chessmen. Contains preliminary section and 58 two-move, 46 three-move, and 8 four-move problems composed by 27 outstanding American problem creators in the last 30 years. Explanation of all terms and exhaustive index. "Just what is wanted for the student," Brian Warley. 112 problems, solutions. vi + 171pp. 5½ x 8. T748 Paperbound \$1.00

CHESS AND CHECKERS: THE WAY TO MASTERSHIP, Edward Lasker. Complete, lucid instructions for the beginner—and valuable suggestions for the advanced player! For both games the great master and teacher presents fundamentals, elementary tactics, and steps toward becoming a superior player. He concentrates on general principles rather than a mass of rules, comprehension rather than brute memory. Historical introduction. 118 diagrams. xiv + 167pp. 5½ x 8. T657 Paperbound \$1.15

WIN AT CHECKERS, M. Hopper. (Formerly CHECKERS). The former World's Unrestricted Checker Champion discusses the principles of the game, expert's shots and traps, problems for the beginner, standard openings, locating your best move, the end game, opening "blitzkrieg" moves, ways to draw when you are behind your opponent, etc. More than 100 detailed questions and answers anticipate your problems. Appendix. 75 problems with solutions and diagrams. Index. 79 figures. xi + 107pp. 5½ x 8. T363 Paperbound \$1.00

GAMES ANCIENT AND ORIENTAL, AND HOW TO PLAY THEM, E. Falkener. A connoisseur's selection of exciting and different games: Oriental varieties of chess, with unusual pieces and moves (including Japanese shogi); the original pachisi; go; reconstructions of lost Roman and Egyptian games; and many more. Full rules and sample games. Now play at home the games that have entertained millions, not on a fad basis, but for millennia. 345 illustrations and

GO AND GO-MOKU: THE ORIENTAL BOARD GAMES, Edward Lasker. Best introduction to Go and its easier sister-game, Go-Moku—games new to Western world, but ancient in China, Japan. Extensively revised work by famed chess master Lasker, Go-player for over 50 years, stresses theory rather than brute memory, presents step-by-step explanation of strategy, gives examples of world championship matches, in game which has replaced chess as favorite of many physicists, mathematicians. 72 diagrams. xix + 215 pp. 5½ x 8.

T613 Paperbound \$1.45

FICTION

FLATLAND, E. A. Abbott. A science-fiction classic of life in a 2-dimensional world that is also a first-rate introduction to such aspects of modern science as relativity and hyperspace. Political, moral, satirical, and humorous overtones have made *FLATLAND* fascinating reading for thousands. 7th edition. New introduction by Banesh Hoffmann. 16 illustrations. 128pp. 5½ x 8.

T1 Paperbound \$1.00

THE WONDERFUL WIZARD OF OZ, L. F. Baum. Only edition in print with all the original W. W. Denslow illustrations in full color—as much a part of “The Wizard” as Tenniel’s drawings are of “Alice in Wonderland.” “The Wizard” is still America’s best-loved fairy tale, in which, as the author expresses it, “The wonderment and joy are retained and the heartaches and nightmares left out.” Now today’s young readers can enjoy every word and wonderful picture of the original book. New introduction by Martin Gardner. A Baum bibliography. 23 full-page color plates. viii + 268pp. 5½ x 8.

T691 Paperbound \$1.45

THE MARVELOUS LAND OF OZ, L. F. Baum. This is the equally enchanting sequel to the “Wizard,” continuing the adventures of the Scarecrow and the Tin Woodman. The hero this time is a little boy named Tip, and all the delightful Oz magic is still present. This is the book with the Animated Saw-horse, the Woggle-Bug, and Jack Pumpkinhead. All the original John R. Neill illustrations, 16 in full color. 287pp. 5½ x 8.

T692 Paperbound \$1.45

FIVE GREAT DOG NOVELS, edited by Blanche Cirkor. The complete original texts of five classic dog novels that have delighted and thrilled millions of children and adults throughout the world with stories of loyalty, adventure, and courage. Full texts of Jack London’s “The Call of the Wild”; John Brown’s “Rab and His Friends”; Alfred Ollivant’s “Bob, Son of Battle”; Marshall Saunders’ “Beautiful Joe”; and Ouida’s “A Dog of Flanders.” 21 illustrations from the original editions. 495pp. 5½ x 8.

T777 Paperbound \$1.50

3 ADVENTURE NOVELS by H. Rider Haggard. Complete texts of “She,” “King Solomon’s Mines,” “Allan Quatermain.” Qualities of discovery; desire for immortality; search for primitive, for what is unadorned by civilization, have kept these novels of African adventure exciting, alive to readers from R. L. Stevenson to George Orwell. 636pp. 5½ x 8.

T584 Paperbound \$2.00

The Space Novels of Jules Verne

TO THE SUN? and OFF ON A COMET!, Jules Verne. Complete texts of two of the most imaginative flights into fancy in world literature display the high adventure that have kept Verne’s novels read for nearly a century. Only unabridged edition of the best translation, by Edward Roth. Large, easily readable type. 50 illustrations selected from first editions. 462pp. 5½ x 8.

T634 Paperbound \$1.75

FROM THE EARTH TO THE MOON and ALL AROUND THE MOON, Jules Verne. Complete editions of two of Verne’s most successful novels, in finest Edward Roth translations, now available after many years out of print. Verne’s visions of submarines, airplanes, television, rockets, interplanetary travel; of scientific and not-so-scientific beliefs; of peculiarities of Americans; all delight and engross us today as much as when they first appeared. Large, easily readable type. 42 illus. from first French edition. 476pp. 5½ x 8.

T633 Paperbound \$1.75

THE CASTING AWAY OF MRS. LECKS AND MRS. ALESHINE, F. R. Stockton. A charming light novel by Frank Stockton, one of America’s finest humorists (and author of “The Lady, or the Tiger?”). This book has made millions of Americans laugh at the reflection of themselves in two middle-aged American women involved in some of the strangest adventures on record. You will laugh, too, as they endure shipwreck, desert island, and blizzard with maddening tranquility. Also contains complete text of “The Dusanter,” sequel to “The Casting Away.” 49 original illustrations by F. D. Steele. vii + 142pp. 5½ x 8.

T743 Paperbound \$1.00

GESTA ROMANORUM, trans. by Charles Swan, ed. by Wynnard Hooper. 181 tales of Greeks, Romans, Britons, Biblical characters, comprise one of greatest medieval story collections, source plots for writers including Shakespeare, Chaucer, Gower, etc. Imaginative tales of wars, incest, thwarted love, magic, fantasy, allegory, humor, tell about kings, prostitutes, philosophers, fair damsels, knights, Noah, pirates, all walks and stations of life. Introduction. Notes. 500pp. 5 1/2" x 8. T535 Paperbound \$1.85

THREE PROPHETIC NOVELS BY H. G. WELLS, edited by E. F. Bleiler. Complete texts of "When the Sleeper Wakes" (1st book printing in 50 years), "A Story of the Days to Come," "The Time Machine" (1st complete printing in book form). Exciting adventures in the future are as enjoyable today as 50 years ago when first printed. Predict TV, movies, intercontinental airplanes, prefabricated houses, air-conditioned cities, etc. First important author to foresee problems of mind control, technological dictatorships. "Absolute best of imaginative fiction," N. Y. Times. Introduction. 335pp. 5 1/2" x 8. T605 Paperbound \$1.45

SEVEN SCIENCE FICTION NOVELS, H. G. Wells. Full unabridged texts of 7 science-fiction novels of the master. Ranging from biology, physics, chemistry, astronomy to sociology and other studies, Mr. Wells extrapolates whole worlds of strange and intriguing character. "One will have to go far to match this for entertainment, excitement, and sheer pleasure. . . ." NEW YORK TIMES. Contents: The Time Machine, The Island of Dr. Moreau, First Men in the Moon, The Invisible Man, The War of the Worlds, The Food of the Gods, In the Days of the Comet. 1015pp. 5 1/2" x 8. T264 Clothbound \$3.95

28 SCIENCE FICTION STORIES OF H. G. WELLS. Two full unabridged novels, MEN LIKE GODS and STAR BEGOTTEN, plus 26 short stories by the master science-fiction writer of all time. Stories of space, time, invention, exploration, future adventure—an indispensable part of the library of everyone interested in science and adventure. PARTIAL CONTENTS: Men Like Gods, The Country of the Blind, In the Abyss, The Crystal Egg, The Man Who Could Work Miracles, A Story of the Days to Come, The Valley of Spiders, and 21 more! 928pp. 5 1/2" x 8. T265 Clothbound \$3.95

DAVID HARUM, E. N. Westcott. This novel of one of the most lovable, humorous characters in American literature is a prime example of regional humor. It continues to delight people who like their humor dry, their characters quaint, and their plots ingenious. First book edition to contain complete novel plus chapter found after author's death. Illustrations from first illustrated edition. 192pp. 5 1/2" x 8. T580 Paperbound \$1.15

HUMOR

THE WIT AND HUMOR OF OSCAR WILDE, ed. by Alvin Redman. Wilde at his most brilliant, in 1000 epigrams exposing weaknesses and hypocrisies of "civilized" society. Divided into 49 categories—sin, wealth, women, America, etc.—to aid writers, speakers. Includes excerpts from his trials, books, plays, criticism. Formerly "The Epigrams of Oscar Wilde." Introduction by Vyvyan Holland, Wilde's only living son. Introductory essay by editor. 260pp. 5 1/2" x 8. T602 Paperbound \$1.00

A NONSENSE ANTHOLOGY, collected by Carolyn Wells. 245 of the best nonsense verses ever written, including nonsense puns, absurd arguments, mock epics and sagas, nonsense ballads, odes, "sick" verses, dog-Latin verses, French nonsense verses, songs. By Edward Lear, Lewis Carroll, Gelett Burgess, W. S. Gilbert, Hilaire Belloc, Peter Newell, Oliver Herford, etc., 83 writers in all plus over four score anonymous nonsense verses. A special section of limericks, plus famous nonsense such as Carroll's "Jabberwocky" and Lear's "The Jumblies" and much excellent verse virtually impossible to locate elsewhere. For 50 years considered the best anthology available. Index of first lines specially prepared for this edition. Introduction by Carolyn Wells. 3 indexes: Title, Author, First lines. xxxiii + 279pp. 5 1/2" x 8. T499 Paperbound \$1.25

THE BAD CHILD'S BOOK OF BEASTS, MORE BEASTS FOR WORSE CHILDREN, and A MORAL ALPHABET, H. Belloc. Hardly an anthology of humorous verse has appeared in the last 50 years without at least a couple of these famous nonsense verses. But one must see the entire volumes—with all the delightful original illustrations by Sir Basil Blackwood—to appreciate fully Belloc's charming and witty verses that play so subacutely on the platitudes of life and morals that beset his day—and ours. A great humor classic. Three books in one. Total of 157pp. 5 1/2" x 8. T749 Paperbound \$1.00

THE DEVIL'S DICTIONARY, Ambrose Bierce. Sardonic and irreverent barbs puncturing the pomposities and absurdities of American politics, business, religion, literature, and arts, by the country's greatest satirist in the classic tradition. Epigrammatic as Shaw, piercing as Swift, American as Mark Twain, Will Rogers, and Fred Allen. Bierce will always remain the favorite of a small coterie of enthusiasts, and of writers and speakers whom he supplies with "some of the most gorgeous witticisms of the English language." (H. L. Mencken) Over 1000 entries in alphabetical order. 144pp. 5 1/2" x 8. T487 Paperbound \$1.00

THE PURPLE COW AND OTHER NONSENSE, Gelett Burgess. The best of Burgess's early nonsense, selected from the first edition of the "Burgess Nonsense Book." Contains many of his most unusual and highly original pieces: 37 nonsense quatrains, the Poems of Patagonia, Alphabet of Famous Goops, and the other hilarious (and rare) adult nonsense that places him in the forefront of American humorists. All pieces are accompanied by the original Burgess illustrations. 123 illustrations. xiii + 113pp. 5½ x 8.

T772 Paperbound \$1.00

THE HUMOROUS VERSE OF LEWIS CARROLL. Almost every poem Carroll ever wrote, the largest collection ever published, including much never published elsewhere: 150 parodies, burlesques, riddles, ballads, acrostics, etc., with 130 original illustrations by Tenniel, Carroll, and others. "Addicts will be grateful . . . there is nothing for the faithful to do but sit down and fall to the banquet," N. Y. Times. Index to first lines. xiv + 446pp. 5 x 8.

T654 Paperbound \$1.85

DIVERSIONS AND DIGRESSIONS OF LEWIS CARROLL. A major new treasure for Carroll fans! Rare privately published humor, fantasy, puzzles, and games by Carroll at his whimsical best, with a new vein of frank satire. Includes many new mathematical amusements and recreations, among them the fragmentary Part III of "Curiosa Mathematica." Contains "The Rectory Umbrella," "The New Belfry," "The Vision of the Three T's," and much more. New 32-page supplement of rare photographs taken by Carroll. x + 375pp. 5½ x 8.

T732 Paperbound \$1.50

THE COMPLETE NONSENSE OF EDWARD LEAR. This is the only complete edition of this master of gentle madness available at a popular price. A BOOK OF NONSENSE, NONSENSE SONGS, MORE NONSENSE SONGS AND STORIES in their entirety with all the old favorites that have delighted children and adults for years. The Dong With A Luminous Nose, The Jumbles, The Owl and the Pussycat, and hundreds of other bits of wonderful nonsense. 214 limericks, 3 sets of Nonsense Botany, 5 Nonsense Alphabets. 546 drawings by Lear himself, and much more. 320pp. 5½ x 8.

T167 Paperbound \$1.00

PECK'S BAD BOY AND HIS PA, George W. Peck. The complete edition, containing both volumes, one of the most widely read of all American humor books. The endless ingenious pranks played by bad boy "Hennerly" on his pa and the grocery man, the outraged pomposity of Pa, the perpetual ridiculing of middle class institutions, are as entertaining today as they were in 1883. No pale sophistications or subtleties, but rather humor vigorous, raw, earthy, imaginative, and, as folk humor often is, sadistic. This peculiarly fascinating book is also valuable to historians and students of American culture as a portrait of an age. 100 original illustrations by True Williams. Introduction by E. F. Bleiler. 347pp. 5½ x 8.

T497 Paperbound \$1.35

FABLES IN SLANG & MORE FABLES IN SLANG, George Ade. 2 complete books of major American humorist in pungent colloquial tradition of Twain, Billings. 1st reprinting in over 30 years includes "The Two Mandolin Players and the Willing Performer," "The Base Ball Fan Who Took the Only Known Cure," "The Slim Girl Who Tried to Keep a Date that was Never Made," 42 other tales of eccentric, perverse, but always funny characters. "Touch of genius," H. L. Mencken. New introduction by E. F. Bleiler. 86 illus. 203pp. 5½ x 8.

T533 Paperbound \$1.00

SINGULAR TRAVELS, CAMPAIGNS, AND ADVENTURES OF BARON MUNCHAUSEN, R. E. Raspe, with 90 illustrations by Gustave Doré. The first edition in over 150 years to reestablish the deeds of the Prince of Liars exactly as Raspe first recorded them in 1785—the genuine Baron Munchausen, one of the most popular personalities in English literature. Included also are the best of the many sequels, written by other hands. Introduction on Raspe by J. Carswell. Bibliography of early editions. xiv + 192pp. 5½ x 8.

T698 Paperbound \$1.00

HOW TO TELL THE BIRDS FROM THE FLOWERS, R. W. Wood. How not to confuse a carrot with a parrot, a grape with an ape, a puffin with nuffin. Delightful drawings, clever puns, absurd little poems point out farfetched resemblances in nature. The author was a leading physicist. Introduction by Margaret Wood White. 106 illus. 60pp. 5½ x 8.

T523 Paperbound 75¢

MATHEMATICS, ELEMENTARY TO INTERMEDIATE

HOW TO CALCULATE QUICKLY, Henry Sticker. This handy volume offers a tried and true method for helping you in the basic mathematics of daily life—addition, subtraction, multiplication, division, fractions, etc. It is designed to awaken your "number sense" or the ability to see relationships between numbers as whole quantities. It is not a collection of tricks working only on special numbers, but a serious course of over 9,000 problems and their solutions, teaching special techniques not taught in schools: left-to-right multiplication, new fast ways of division, etc. 5 or 10 minutes daily use will double or triple your calculation speed. Excellent for the scientific worker who is at home in higher math, but is not satisfied with his speed and accuracy in lower mathematics. 256pp. 5 x 7¼.

T295 Paperbound \$1.00

TEACH YOURSELF books. For adult self-study, for refresher and supplementary study.

The most effective series of home study mathematics books on the market! With absolutely no outside help, they will teach you as much as any similar college or high-school course, or will helpfully supplement any such course. Each step leads directly to the next, each question is anticipated. Numerous lucid examples and carefully-wrought practice problems illustrate meanings. Not skimpy outlines, not surveys, not usual classroom texts, these 204- to 380-page books are packed with the finest instruction you'll find anywhere for adult self-study.

TEACH YOURSELF ALGEBRA, P. Abbott. Formulas, coordinates, factors, graphs of quadratic functions, quadratic equations, logarithms, ratio, irrational numbers, arithmetical, geometrical series, much more. 1241 problems, solutions. Tables. 52 illus. 307pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$.
Clothbound **\$2.00**

TEACH YOURSELF GEOMETRY, P. Abbott. Solids, lines, points, surfaces, angle measurement, triangles, theorem of Pythagoras, polygons, loci, the circle, tangents, symmetry, solid geometry, prisms, pyramids, solids of revolution, etc. 343 problems, solutions. 268 illus. 334pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$.
Clothbound **\$2.00**

TEACH YOURSELF TRIGONOMETRY, P. Abbott. Geometrical foundations, indices, logarithms, trigonometrical ratios, relations between sides, angles of triangle, circular measure, trig. ratios of angles of any magnitude, much more. Requires elementary algebra, geometry. 465 problems, solutions. Tables. 102 illus. 204pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$.
Clothbound **\$2.00**

TEACH YOURSELF THE CALCULUS, P. Abbott. Variations in functions, differentiation, solids of revolution, series, elementary differential equations, areas by integral calculus, much more. Requires algebra, trigonometry. 970 problems, solutions. Tables. 89 illus. 380pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$.
Clothbound **\$2.00**

TEACH YOURSELF THE SLIDE RULE, B. Snodgrass. Fractions, decimals, A-D scales, log-log scales, trigonometrical scales, indices, logarithms. Commercial, precision, electrical, dualistic, Brighton rules. 80 problems, solutions. 10 illus. 207pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$.
Clothbound **\$2.00**

See also: **TEACH YOURSELF ELECTRICITY, C. W. Wilman; TEACH YOURSELF HEAT ENGINES, E. De Ville; TEACH YOURSELF MECHANICS, P. Abbott.**



HOW DO YOU USE A SLIDE RULE? by A. A. Merrill. Not a manual for mathematicians and engineers, but a lucid step-by-step explanation that presents the fundamental rules clearly enough to be understood by anyone who could benefit by the use of a slide rule in his work or business. This work concentrates on the 2 most important operations: multiplication and division. 10 easy lessons, each with a clear drawing, will save you countless hours in your banking, business, statistical, and other work. First publication. Index. 2 Appendices. 10 illustrations. 78 problems, all with answers. vi + 36pp. 6 $\frac{7}{8}$ x 9 $\frac{1}{4}$.
T62 Paperbound **60¢**

THEORY OF OPERATION OF THE SLIDE RULE, J. P. Ellis. Not a skimpy "instruction manual", but an exhaustive treatment that will save you uncounted hours throughout your career. Supplies full understanding of every scale on the Log Log Duplex Decitrig type of slide rule. Shows the most time-saving methods, and provides practice useful in the widest variety of actual engineering situations. Each operation introduced in terms of underlying logarithmic theory. Summary of prerequisite math. First publication. Index. 198 figures. Over 450 problems with answers. Bibliography. 12 Appendices. ix + 289pp. 5 $\frac{3}{8}$ x 8.
S727 Paperbound **\$1.50**

ARITHMETICAL EXCURSIONS: AN ENRICHMENT OF ELEMENTARY MATHEMATICS, H. Bowers and J. Bowers. For students who want unusual methods of arithmetic never taught in school; for adults who want to increase their number sense. Little known facts about the most simple numbers, arithmetical entertainments and puzzles, figurate numbers, number chains, mysteries and folklore of numbers, the "Hin-dog-abc" number system, etc. First publication. Index. 529 numbered problems and diversions, all with answers. Bibliography. 50 figures. xiv + 320pp. 5 $\frac{3}{8}$ x 8.
T770 Paperbound **\$1.65**

APPLIED MATHEMATICS FOR RADIO AND COMMUNICATIONS ENGINEERS, C. E. Smith. No extraneous material here!—only the theories, equations, and operations essential and immediately useful for radio work. Can be used as refresher, as handbook of applications and tables, or as full home-study course. Ranges from simplest arithmetic through calculus, series, and wave forms, hyperbolic trigonometry, simultaneous equations in mesh circuits, etc. Supplies applications right along with each math topic discussed. 22 useful tables of functions, formulas, logs, etc. Index. 166 exercises, 140 examples, all with answers. 95 diagrams. Bibliography. x + 336pp. 5 $\frac{3}{8}$ x 8.
S141 Paperbound **\$1.75**

HIGHER MATHEMATICS FOR STUDENTS OF CHEMISTRY AND PHYSICS, J. W. Mellor. Not abstract, but practical, building its problems out of familiar laboratory material, this covers differential calculus, coordinate, analytical geometry, functions, integral calculus, infinite series, numerical equations, differential equations, Fourier's theorem, probability, theory of errors, calculus of variations, determinants. "If the reader is not familiar with this book, it will repay him to examine it," CHEM. & ENGINEERING NEWS. 800 problems. 189 figures. Bibliography. xxi + 641pp. 5% x 8. S193 Paperbound \$2.25

TRIGONOMETRY REFRESHER FOR TECHNICAL MEN, A. Albert Klaf. 913 detailed questions and answers cover the most important aspects of plane and spherical trigonometry. They will help you to brush up or to clear up difficulties in special areas. The first portion of this book covers plane trigonometry, including angles, quadrants, trigonometrical functions, graphical representation, interpolation, equations, logarithms, solution of triangle, use of the slide rule and similar topics. 188 pages then discuss application of plane trigonometry to special problems in navigation, surveying, elasticity, architecture, and various fields of engineering. Small angles, periodic functions, vectors, polar coordinates, de Moivre's theorem are fully examined. The third section of the book then discusses spherical trigonometry and the solution of spherical triangles, with their applications to terrestrial and astronomical problems. Methods of saving time with numerical calculations, simplification of principal functions of angle, much practical information make this a most useful book. 913 questions answered. 1738 problems, answers to odd numbers. 494 figures. 24 pages of useful formulae, functions. Index. x + 629pp. 5% x 8. T371 Paperbound \$2.00

CALCULUS REFRESHER FOR TECHNICAL MEN, A. Albert Klaf. This book is unique in English as a refresher for engineers, technicians, students who either wish to brush up their calculus or to clear up uncertainties. It is not an ordinary text, but an examination of most important aspects of integral and differential calculus in terms of the 756 questions most likely to occur to the technical reader. The first part of this book covers simple differential calculus, with constants, variables, functions, increments, derivatives, differentiation, logarithms, curvature of curves, and similar topics. The second part covers fundamental ideas of integration, inspection, substitution, transformation, reduction, areas and volumes, mean value, successive and partial integration, double and triple integration. Practical aspects are stressed rather than theoretical. A 50-page section illustrates the application of calculus to specific problems of civil and nautical engineering, electricity, stress and strain, elasticity, industrial engineering, and similar fields.—756 questions answered. 566 problems, mostly answered. 36 pages of useful constants, formulae for ready reference. Index. v + 431pp. 5% x 8. T370 Paperbound \$2.00

TEXTBOOK OF ALGEBRA, G. Chrystal. One of the great mathematical textbooks, still about the best source for complete treatments of the topics of elementary algebra; a chief reference work for teachers and students of algebra in advanced high school and university courses, or for the mathematician working on problems of elementary algebra or looking for a background to more advanced topics. Ranges from basic laws and processes to extensive examination of such topics as limits, infinite series, general properties of integral numbers, and probability theory. Emphasis is on algebraic form, the foundation of analytical geometry and the key to modern developments in algebra. Prior course in algebra is desirable, but not absolutely necessary. Includes theory of quotients, distribution of products, arithmetical theory of surds, theory of interest, permutations and combinations, general expansion theorems, recurring fractions, and much, much more. Two volume set. Index in each volume. Over 1500 exercises, approximately half with answers. Total of xlviii + 1187pp. 5% x 8.

S750 Vol I Paperbound \$2.35

S751 Vol II Paperbound \$2.35

The set \$4.70

COLLEGE ALGEBRA, H. B. Fine. Standard college text that gives a systematic and deductive structure to algebra; comprehensive, connected, with emphasis on theory. Discusses the commutative, associative, and distributive laws of number in unusual detail, and goes on with undetermined coefficients, quadratic equations, progressions, logarithms, permutations, probability, power series, and much more. Still most valuable elementary-intermediate text on the science and structure of algebra. Index. 1560 problems, all with answers. x + 631pp. 5% x 8. T211 Paperbound \$2.00

THE CONTINUUM AND OTHER TYPES OF SERIAL ORDER, E. V. Huntington. This famous book gives a systematic elementary account of the modern theory of the continuum as a type of serial order. Based on the Cantor-Dedekind ordinal theory, which requires no technical knowledge of higher mathematics, it offers an easily followed analysis of ordered classes, discrete and dense series, continuous series, Cantor's transfinite numbers. 2nd edition. Index. viii + 82pp. 5% x 8. S129 Clothbound \$2.75

S130 Paperbound \$1.00

A TREATISE ON PLANE AND ADVANCED TRIGONOMETRY, E. W. Hobson. Extraordinarily wide coverage, going beyond usual college level trig, one of the few works covering advanced trig in full detail. By a great expositor with unerring anticipation and lucid clarification of potentially difficult points. Includes circular functions; expansion of functions of multiple angle; trig tables; relations between sides and angles of triangle; complex numbers; etc. Many problems solved completely. "The best work on the subject." Nature. Formerly entitled "A Treatise on Plane Trigonometry." 689 examples. 6 figures. xvi + 383pp. 5% x 8. S353 Paperbound \$1.95

FAMOUS PROBLEMS OF ELEMENTARY GEOMETRY, Felix Klein. Expanded version of the 1894 Easter lectures at Göttingen. 3 problems of classical geometry, in an excellent mathematical treatment by a famous mathematician: squaring the circle, trisecting angle, doubling cube. Considered with full modern implications: transcendental numbers, π , etc. Notes by R. Archibald. 16 figures. xi + 92pp. 5% x 8. T348 Clothbound \$1.50
T298 Paperbound \$1.00



ELEMENTARY MATHEMATICS FROM AN ADVANCED STANDPOINT, Felix Klein.

This classic text is an outgrowth of Klein's famous integration and survey course at Göttingen. Using one field of mathematics to interpret, adjust, illuminate another, it covers basic topics in each area, illustrating its discussion with extensive analysis. It is especially valuable in considering areas of modern mathematics. "Makes the reader feel the inspiration of . . . a great mathematician, inspiring teacher . . . with deep insight into the foundations and interrelations," BULLETIN, AMERICAN MATHEMATICAL SOCIETY.

Vol. 1. ARITHMETIC, ALGEBRA, ANALYSIS. Introducing the concept of function immediately, it enlivens abstract discussion with graphical and geometrically perceptual methods. Partial contents: natural numbers, extension of the notion of number, special properties, complex numbers. Real equations with real unknowns, complex quantities. Logarithmic, exponential functions, goniometric functions, infinitesimal calculus. Transcendence of e and π , theory of assemblages. Index. 125 figures. ix + 274pp. 5% x 8. S150 Paperbound \$1.75

Vol. 2. GEOMETRY. A comprehensive view which accompanies the space perception inherent in geometry with analytic formulas which facilitate precise formulation. Partial contents: Simplest geometric manifolds: line segment, Grassmann determinant principles, classification of configurations of space, derivative manifolds. Geometric transformations: affine transformations, projective, higher point transformations, theory of the imaginary. Systematic discussion of geometry and its foundations. Indexes. 141 illustrations. ix + 214pp. 5% x 8. S151 Paperbound \$1.75

* * *

COORDINATE GEOMETRY, L. P. Eisenhart. Thorough, unified introduction. Unusual for advancing in dimension within each topic (treats together circle, sphere; polar coordinates, 3-dimensional coordinate systems; conic sections, quadric surfaces), affording exceptional insight into subject. Extensive use made of determinants, though no previous knowledge of them is assumed. Algebraic equations of 1st degree, 2 and 3 unknowns, carried further than usual in algebra courses. Over 500 exercises. Introduction. Appendix. Index. Bibliography. 43 illustrations. 310pp. 5% x 8. S600 Paperbound \$1.65

MONOGRAPHS ON TOPICS OF MODERN MATHEMATICS, edited by J. W. A. Young. Advanced mathematics for persons who haven't gone beyond or have forgotten high school algebra. 9 monographs on foundation of geometry, modern pure geometry, non-Euclidean geometry, fundamental propositions of algebra, algebraic equations, functions, calculus, theory of numbers, etc. Each monograph gives proofs of important results, and descriptions of leading methods, to provide wide coverage. New Introduction by Prof. M. Kline, N. Y. University. 100 diagrams. xvi + 416pp. 6% x 9% . S289 Paperbound \$2.00

MATHEMATICS, INTERMEDIATE TO ADVANCED

Geometry

THE FOUNDATIONS OF EUCLIDEAN GEOMETRY, H. G. Forder. The first rigorous account of Euclidean geometry, establishing propositions without recourse to empiricism, and without multiplying hypotheses. Corrects many traditional weaknesses of Euclidean proofs, and investigates the problems imposed on the axiom system by the discoveries of Bolya and Lobachevsky. Some topics discussed are Classes and Relations; Axioms for Magnitudes; Congruence and Similarity; Algebra of Points; Hessenberg's Theorem; Continuity; Existence of Parallels; Reflections; Rotations; Isometries; etc. Invaluable for the light it throws on foundations of math. Lists: Axioms employed, Symbols, Constructions. 295pp. 5% x 8. S481 Paperbound \$2.00

ADVANCED EUCLIDEAN GEOMETRY, R. A. Johnson. For years the standard textbook on advanced Euclidean geometry, requires only high school geometry and trigonometry. Explores in unusual detail and gives proofs of hundreds of relatively recent theorems and corollaries, many formerly available only in widely scattered journals. Covers tangent circles, the theorem of Miquel, symmedian point, pedal triangles and circles, the Brocard configuration, and much more. Formerly "Modern Geometry." Index. 107 diagrams. xiii + 319pp. 5% x 8. S669 Paperbound \$1.65

NON-EUCLIDEAN GEOMETRY, Roberto Bonola. The standard coverage of non-Euclidean geometry. It examines from both a historical and mathematical point of view the geometries which have arisen from a study of Euclid's 5th postulate upon parallel lines. Also included are complete texts, translated, of Bolyai's **THEORY OF ABSOLUTE SPACE**, Lobachevsky's **THEORY OF PARALLELS**. 180 diagrams. 431pp. 5% x 8. S27 Paperbound \$1.95

ELEMENTS OF NON-EUCLIDEAN GEOMETRY, D. M. Y. Sommerville. Unique in proceeding step-by-step, in the manner of traditional geometry. Enables the student with only a good knowledge of high school algebra and geometry to grasp elementary hyperbolic, elliptic, analytic non-Euclidean geometries; space curvature and its philosophical implications; theory of radical axes; homothetic centres and systems of circles; parataxy and parallelism; absolute measure; Gauss' proof of the defect area theorem; geodesic representation; much more, all with exceptional clarity. 126 problems at chapter endings provide progressive practice and familiarity. 133 figures. Index. xvi + 274pp. 5% x 8. S460 Paperbound \$1.50

HIGHER GEOMETRY: AN INTRODUCTION TO ADVANCED METHODS IN ANALYTIC GEOMETRY, F. S. Woods. Exceptionally thorough study of concepts and methods of advanced algebraic geometry (as distinguished from differential geometry). Exhaustive treatment of 1-, 2-, 3-, and 4-dimensional coordinate systems, leading to n-dimensional geometry in an abstract sense. Covers projectivity, tetracyclic coordinates, contact transformation, pentaspherical coordinates, much more. Based on M.I.T. lectures, requires sound preparation in analytic geometry and some knowledge of determinants. Index. Over 350 exercises. References. 60 figures. x + 423pp. 5% x 8. S737 Paperbound \$2.00

ELEMENTS OF PROJECTIVE GEOMETRY, L. Cremona. Outstanding complete treatment of projective geometry by one of the foremost 19th century geometers. Detailed proofs of all fundamental principles, stress placed on the constructive aspects. Covers homology, law of duality, anharmonic ratios, theorems of Pascal and Brianchon, foci, polar reciprocal figures, etc. Only ordinary geometry necessary to understand this honored classic. Index. Over 150 fully worked out examples and problems. 252 diagrams. xx + 302pp. 5% x 8. S668 Paperbound \$1.75

A TREATISE ON THE DIFFERENTIAL GEOMETRY OF CURVES AND SURFACES, L. P. Eisenhart. Introductory treatise especially for the graduate student, for years a highly successful textbook. More detailed and concrete in approach than most more recent books. Covers space curves, osculating planes, moving axes, Gauss' method, the moving trihedral, geodesics, conformal representation, etc. Last section deals with deformation of surfaces, rectilinear congruences, cyclic systems, etc. Index. 683 problems. 30 diagrams. xii + 474pp. 5% x 8. S667 Paperbound \$2.75

A TREATISE ON ALGEBRAIC PLANE CURVES, J. L. Coolidge. Unabridged reprinting of one of few full coverages in English, offering detailed introduction to theory of algebraic plane curves and their relations to geometry and analysis. Treats topological properties, Riemann-Roch theorem, all aspects of wide variety of curves including real, covariant, polar, containing series of a given sort, elliptic, polygonal, rational, the pencil, two parameter nets, etc. This volume will enable the reader to appreciate the symbolic notation of Aronhold and Clebsch. Bibliography. Index. 17 illustrations. xxiv + 513pp. 5% x 8. S543 Paperbound \$2.45

AN INTRODUCTION TO THE GEOMETRY OF N DIMENSIONS, D. M. Y. Sommerville. An introduction presupposing no prior knowledge of the field, the only book in English devoted exclusively to higher dimensional geometry. Discusses fundamental ideas of incidence, parallelism, perpendicularity, angles between linear space; enumerative geometry; analytical geometry from projective and metric points of view; polytopes; elementary ideas in analysis situs; content of hyper-spacial figures. Bibliography. Index. 60 diagrams. 196pp. 5% x 8. S494 Paperbound \$1.50

GEOMETRY OF FOUR DIMENSIONS, H. P. Manning. Unique in English as a clear, concise introduction. Treatment is synthetic, and mostly Euclidean, although in hyperplanes and hyperspheres at infinity, non-Euclidean geometry is used. Historical introduction. Foundations of 4-dimensional geometry. Perpendicularity, simple angles. Angles of planes, higher order. Symmetry, order, motion; hyperpyramids, hypercones, hyperspheres; figures with parallel elements; volume, hypervolume in space; regular polyhedroids. Glossary. 78 figures. ix + 348pp. 5% x 8. S182 Paperbound \$1.95

ELEMENTARY CONCEPTS OF TOPOLOGY, P. Alexandroff. First English translation of the famous brief introduction to topology for the beginner or for the mathematician not undertaking extensive study. This unusually useful intuitive approach deals primarily with the concepts of complex, cycle, and homology, and is wholly consistent with current investigations. Ranges from basic concepts of set-theoretic topology to the concept of Betti groups. "Glowing example of harmony between intuition and thought," David Hilbert. Translated by A. E. Farley. Introduction by D. Hilbert. Index. 25 figures. 73pp. 5% x 8. S747 Paperbound \$1.00

THE WORKS OF ARCHIMEDES, edited by T. L. Heath. All the known works of the great Greek mathematician are contained in this one volume, including the recently discovered Method of Archimedes. Contains: On Sphere & Cylinder, Measurement of a Circle, Spirals, Conoids, Spheroids, etc. This is the definitive edition of the greatest mathematical intellect of the ancient world. 186-page study by Heath discusses Archimedes and the history of Greek mathematics. Bibliography. 563pp. 5% x 8. S9 Paperbound \$2.00

THE THIRTEEN BOOKS OF EUCLID'S ELEMENTS, edited by Sir Thomas Heath. Definitive edition of one of the very greatest classics of Western world. Complete English translation of Heiberg text, together with spurious Book XIV. Detailed 150-page introduction discussing aspects of Greek and Medieval mathematics. Euclid, texts, commentators, etc. Paralleling the text is an elaborate critical apparatus analyzing each definition, proposition, postulate, covering textual matters, mathematical analysis, commentators of all times, refutations, supports, extrapolations, etc. This is the FULL EUCLID. Unabridged reproduction of Cambridge U. 2nd edition. 3 volumes. Total of 995 figures, 1426pp. 5 3/8 x 8.

\$88,89,90, 3 volume set, paperback \$6.00

THE GEOMETRY OF RENE DESCARTES. With this book Descartes founded analytical geometry. Excellent Smith-Latham translation, plus original French text with Descartes' own diagrams. Contains Problems the Construction of Which Requires Only Straight Lines and Circles; On the Nature of Curved Lines; On the Construction of Solid or Supersolid Problems. Notes. Diagrams. 258pp. 5 1/2 x 8.

\$68 Paperbound \$1.50

See also: **FOUNDATIONS OF GEOMETRY**, B. Russell; **THE PHILOSOPHY OF SPACE AND TIME**, H. Reichenbach; **FAMOUS PROBLEMS OF ELEMENTARY GEOMETRY**, F. Klein; **MONOGRAPHS ON TOPICS OF MODERN MATHEMATICS**, ed. by J. W. Young.

Calculus and function theory, Fourier theory, real and complex functions, determinants

A COLLECTION OF MODERN MATHEMATICAL CLASSICS, edited by R. Bellman. 13 classic papers, complete in their original languages, by Hermite, Hardy and Littlewood, Tchebychev, Fejér, Fredholm, Fuchs, Hurwitz, Weyl, van der Pol, Birkhoff, Kellogg, von Neumann, and Hilbert. Each of these papers, collected here for the first time, triggered a burst of mathematical activity, providing useful new generalizations or stimulating fresh investigations. Topics discussed include classical analysis, periodic and almost periodic functions, analysis and number theory, integral equations, theory of approximation, non-linear differential equations, and functional analysis. Brief introductions and bibliographies to each paper. xii + 292pp. 6 x 9.

\$730 Paperbound \$2.00

MATHEMATICS OF MODERN ENGINEERING, E. G. Keller and R. E. Doherty. Written for the Advanced Course in Engineering of the General Electric Corporation, deals with the engineering use of determinants, tensors, the Heaviside operational calculus, dyadics, the calculus of variations, etc. Presents underlying principles fully, but purpose is to teach engineers to deal with modern engineering problems, and emphasis is on the perennial engineering attack of set-up and solve. Indexes. Over 185 figures and tables. Hundreds of exercises, problems, and worked-out examples. References. Two volume set. Total of xxxiii + 623pp. 5 1/2 x 8.

\$734 Vol I Paperbound \$1.65

\$735 Vol II Paperbound \$1.65

The set \$3.30

MATHEMATICAL METHODS FOR SCIENTISTS AND ENGINEERS, L. P. Smith. For scientists and engineers, as well as advanced math students. Full investigation of methods and practical description of conditions under which each should be used. Elements of real functions, differential and integral calculus, space geometry, theory of residues, vector and tensor analysis, series of Bessel functions, etc. Each method illustrated by completely-worked-out examples, mostly from scientific literature. 368 graded unsolved problems. 100 diagrams. x + 453pp. 5 1/2 x 8 3/8.

\$220 Paperbound \$2.00

THEORY OF FUNCTIONS AS APPLIED TO ENGINEERING PROBLEMS, edited by R. Rothe, F. Ollendorff, and K. Pohlhausen. A series of lectures given at the Berlin Institute of Technology that shows the specific applications of function theory in electrical and allied fields of engineering. Six lectures provide the elements of function theory in a simple and practical form, covering complex quantities and variables, integration in the complex plane, residue theorems, etc. Then 5 lectures show the exact uses of this powerful mathematical tool, with full discussions of problem methods. Index. Bibliography. 108 figures. x + 189pp. 5 1/2 x 8.

\$733 Paperbound \$1.35

ADVANCED CALCULUS, E. B. Wilson. An unabridged reprinting of the work which continues to be recognized as one of the most comprehensive and useful texts in the field. It contains an immense amount of well-presented, fundamental material, including chapters on vector functions, ordinary differential equations, special functions, calculus of variations, etc., which are excellent introductions to these areas. For students with only one year of calculus, more than 1300 exercises cover both pure math and applications to engineering and physical problems. For engineers, physicists, etc., this work, with its 54 page introductory review, is the ideal reference and refresher. Index. ix + 566pp. 5 1/2 x 8.

\$504 Paperbound \$2.45

CALCULUS OF VARIATIONS, A. R. Forsyth. Methods, solutions, rather than determination of weakest valid hypotheses. Over 150 examples completely worked-out show use of Euler, Legendre, Jacobi, Weierstrass tests for maxima, minima. Integrals with one original dependent variable; with derivatives of 2nd order; two dependent variables, one independent variable; double integrals involving 1 dependent variable, 2 first derivatives; double integrals involving partial derivatives of 2nd order; triple integrals; much more. 50 diagrams. 678pp. 5% x 8%.

S622 Paperbound \$2.95

LECTURES ON THE CALCULUS OF VARIATIONS, O. Bolza. Analyzes in detail the fundamental concepts of the calculus of variations, as developed from Euler to Hilbert, with sharp formulations of the problems and rigorous demonstrations of their solutions. More than a score of solved examples; systematic references for each theorem. Covers the necessary and sufficient conditions; the contributions made by Euler, Du Bois Reymond, Hilbert, Weierstrass, Legendre, Jacobi, Erdmann, Kneser, and Gauss; and much more. Index. Bibliography. xi + 271pp. 5% x 8.

S218 Paperbound \$

A TREATISE ON THE CALCULUS OF FINITE DIFFERENCES, G. Boole. A classic in the literature of the calculus. Thorough, clear discussion of basic principles, theorems, methods. Covers MacLaurin's and Herschel's theorems, mechanical quadrature, factorials, periodical constants, Bernoulli's numbers, difference-equations (linear, mixed, and partial), etc. Stresses analogies with differential calculus. 236 problems, answers to the numerical ones. viii + 336pp. 5% x 8.

S695 Paperbound \$1.85

THE ANALYTICAL THEORY OF HEAT, Joseph Fourier. This book, which revolutionized mathematical physics, is listed in the Great Books program, and many other listings of great books. It has been used with profit by generations of mathematicians and physicists who are interested in either heat or in the application of the Fourier integral. Covers cause and reflection of rays of heat, radiant heating, heating of closed spaces, use of trigonometric series in the theory of heat, Fourier integral, etc. Translated by Alexander Freeman. 20 figures. xxii + 466pp. 5% x 8.

S93 Paperbound \$2.00

AN INTRODUCTION TO FOURIER METHODS AND THE LAPLACE TRANSFORMATION, Philip Franklin. Concentrates upon essentials, enabling the reader with only a working knowledge of calculus to gain an understanding of Fourier methods in a broad sense, suitable for most applications. This work covers complex qualities with methods of computing elementary functions for complex values of the argument and finding approximations by the use of charts; Fourier series and integrals with half-range and complex Fourier series; harmonic analysis; Fourier and Laplace transformations, etc.; partial differential equations with applications to transmission of electricity, etc. The methods developed are related to physical problems of heat flow, vibrations, electrical transmission, electromagnetic radiation, etc. 828 problems with answers. Formerly entitled "Fourier Methods." Bibliography. Index. x + 289pp. 5% x 8.

S452 Paperbound \$1.75

THE FOURIER INTEGRAL AND CERTAIN OF ITS APPLICATIONS, Norbert Wiener. The only book-length study of the Fourier integral as link between pure and applied math. An expansion of lectures given at Cambridge. Partial contents: Plancherel's theorem, general Tauberian theorem, special Tauberian theorems, generalized harmonic analysis. Bibliography. viii + 201pp. 5% x 8.

S272 Paperbound \$1.50

INTRODUCTION TO THE THEORY OF FOURIER'S SERIES AND INTEGRALS, H. S. Carslaw. 3rd revised edition. This excellent introduction is an outgrowth of the author's courses at Cambridge. Historical introduction, rational and irrational numbers, infinite sequences and similar topics. Appendixes discuss practical harmonic analysis, periodogram analysis. Lebesgues and theory. Indexes. 84 examples, bibliography. xiii + 368pp. 5% x 8.

S48 Paperbound \$2.00

FOURIER'S SERIES AND SPHERICAL HARMONICS, W. E. Byerly. Continues to be recognized as one of most practical, useful expositions. Functions, series, and their differential equations are concretely explained in great detail; theory is applied constantly to practical problems, which are fully and lucidly worked out. Appendix includes 6 tables of surface zonal harmonics, hyperbolic functions, Bessel's functions. Bibliography. 190 problems, approximately half with answers. ix + 287pp. 5% x 8.

S536 Paperbound \$1.75

ASYMPTOTIC EXPANSIONS, A. Erdélyi. The only modern work available in English, this is an unabridged reproduction of a monograph prepared for the Office of Naval Research. It discusses various procedures for asymptotic evaluation of integrals containing a large parameter and solutions of ordinary linear differential equations. Bibliography of 71 items. vi + 108pp. 5% x 8.

S318 Paperbound \$1.35

LINEAR INTEGRAL EQUATIONS, W. V. Lovitt. Systematic survey of general theory, with some application to differential equations, calculus of variations, problems of math, physics. Partial contents: integral equation of 2nd kind by successive substitutions; Fredholm's equation as ratio of 2 integral series in lambda, applications of the Fredholm theory, Hilbert-Schmidt theory of symmetric kernels, application, etc. Neumann, Dirichlet, vibratory problems. Index. ix + 253pp. 5% x 8.

S175 Clothbound \$3.50

S176 Paperbound \$1.60

THEORY OF FUNCTIONALS AND OF INTEGRAL AND INTEGRO-DIFFERENTIAL EQUATIONS, Vito Volterra. Unabridged republication of the only English translation. An exposition of the general theory of the functions depending on a continuous set of values of another function, based on the author's fundamental notion of the transition from a finite number of variables to a continually infinite number. Though dealing primarily with integral equations, much material on calculus of variations is included. The work makes no assumption of previous knowledge on the part of the reader. It begins with fundamental material and proceeds to Generalization of Analytic Functions, Integro-Differential Equations, Functional Derivative Equations, Applications, Other Directions of Theory of Functionals, etc. New introduction by G. C. Evans. Bibliography and criticism of Volterra's work by E. Whittaker. Bibliography. Index of authors cited. Index of subjects. xxxx + 226pp. 5½ x 8. S502 Paperbound \$1.75

AN ELEMENTARY TREATISE ON ELLIPTIC FUNCTIONS, A. Cayley. Still the fullest and clearest text on the theories of Jacobi and Legendre for the advanced student (and an excellent supplement for the beginner). A masterpiece of exposition by the great 19th century British mathematician (creator of the theory of matrices and abstract geometry), it covers the addition-theory, Landen's theorem, the 3 kinds of elliptic integrals, transformations, the q-functions, reduction of a differential expression, and much more. Index. xii + 386pp. 5½ x 8. S728 Paperbound \$2.00

THE APPLICATIONS OF ELLIPTIC FUNCTIONS, A. G. Greenhill. Modern books forgo detail for sake of brevity—this book offers complete exposition necessary for proper understanding, use of elliptic integrals. Formulas developed from definite physical, geometric problems; examples representative enough to offer basic information in widely useable form. Elliptic integrals, addition theorem, algebraical form of addition theorem, elliptic integrals of 2nd, 3rd kind, double periodicity, resolution into factors, series, transformation, etc. Introduction. Index. 25 illus. xi + 357pp. 5½ x 8. S603 Paperbound \$1.75

THE THEORY OF FUNCTIONS OF REAL VARIABLES, James Pierpont. A 2-volume authoritative exposition, by one of the foremost mathematicians of his time. Each theorem stated with all conditions, then followed by proof. No need to go through complicated reasoning to discover conditions added without specific mention. Includes a particularly complete, rigorous presentation of theory of measure; and Pierpont's own work on a theory of Lebesgue integrals, and treatment of area of a curved surface. Partial contents, Vol. 1: rational numbers, exponentials, logarithms, point aggregates, maxima, minima, proper integrals, improper integrals, multiple proper integrals, continuity, discontinuity, indeterminate forms. Vol. 2: point sets, proper integrals, series, power series, aggregates, ordinal numbers, discontinuous functions, sub-, infra-uniform convergence, much more. Index. 95 illustrations. 1229pp. 5½ x 8. S558-9, 2 volume set, paperbound \$4.90

FUNCTIONS OF A COMPLEX VARIABLE, James Pierpont. Long one of best in the field. A thorough treatment of fundamental elements, concepts, theorems. A complete study, rigorous, detailed, with carefully selected problems worked out to illustrate each topic. Partial contents: arithmetical operations, real term series, positive term series, exponential functions, integration, analytic functions, asymptotic expansions, functions of Weierstrass, Legendre, etc. Index. List of symbols. 122 illus. 597pp. 5½ x 8. S560 Paperbound \$2.45

ELEMENTS OF THE THEORY OF REAL FUNCTIONS, J. E. Littlewood. Based on lectures given at Trinity College, Cambridge, this book has proved to be extremely successful in introducing graduate students to the modern theory of functions. It offers a full and concise coverage of classes and cardinal numbers, well-ordered series, other types of series, and elements of the theory of sets of points. 3rd revised edition. vii + 71pp. 5½ x 8.

S171 Clothbound \$2.85
S172 Paperbound \$1.25

TRANSCENDENTAL AND ALGEBRAIC NUMBERS, A. O. Gelfond. First English translation of work by leading Soviet mathematician. Thue-Siegel theorem, its p-adic analogue, on approximation of algebraic numbers by numbers in fixed algebraic field; Hermite-Lindemann theorem on transcendency of Bessel functions, solutions of other differential equations; Gelfond-Schneider theorem on transcendency of α^b to power β ; Schneider's work on elliptic functions, with method developed by Gelfond. Translated by L. F. Boron. Index. Bibliography. 200pp. 5½ x 8. S615 Paperbound \$1.75

Dover publishes books on art, music, philosophy, literature, languages, history, social sciences, psychology, handicrafts, orientalia, puzzles and entertainments, chess, pets and gardens, books explaining science, intermediate and higher mathematics mathematical physics, engineering, biological sciences, earth sciences, classics of science, etc. Write to:

*Dept. catrr.
Dover Publications, Inc.
180 Varil Street, New York, N.Y.*